

# From Vlasov-Poisson to Korteweg-de Vries and Zakharov-Kuznetsov

Daniel Han-Kwan

École Normale Supérieure, Département de Mathématiques et Applications,  
45 rue d'Ulm, 75230 Paris Cedex 05 France.  
E-mail: daniel.han-kwan@ens.fr

The date of receipt and acceptance will be inserted by the editor

**Abstract:** We introduce a long wave scaling for the Vlasov-Poisson equation and derive, in the cold ions limit, the Korteweg-De Vries equation (in 1D) and the Zakharov-Kuznetsov equation (in higher dimensions, in the presence of an external magnetic field). The proofs are based on the relative entropy method.

## 1. The long wave scaling of the Vlasov-Poisson equation

*1.1. The Vlasov-Poisson system for ions with small mass electrons.* We consider the Vlasov-Poisson system which describes the evolution of ions in a plasma. We assume that due to their small mass, electrons in the plasma instantaneously reach their thermodynamic equilibrium, so that their density  $n_e$  follows the Maxwell-Boltzmann law:

$$n_e = e^{\phi/T_e}, \quad (1.1)$$

denoting by  $\phi$  the electric potential and  $T_e > 0$  the scaled temperature of electrons, taken equal to 1 in the following. We then obtain the Vlasov-Poisson system for ions (where  $t \geq 0$ ,  $x \in \mathbb{T}^d$  or  $\mathbb{R}^d$ ,  $v \in \mathbb{R}^d$ ,  $d = 1, 2, 3$ ):

$$\begin{cases} \partial_t f + v \cdot \nabla_x f + (E + v \wedge b) \cdot \nabla_v f = 0, \\ E = -\nabla_x \phi, \\ -\Delta_x \phi + e^\phi = \int_{\mathbb{R}^d} f dv, \\ f|_{t=0} = f_0. \end{cases} \quad (1.2)$$

We refer to [26] for a more complete discussion on this system. In these equations,  $f(t, x, v)$  stands for the *distribution function* of the ions, which allows to describe their statistical distribution: this means that  $f(t, x, v) dx dv$  expresses the density of ions, at time  $t$ , whose position is close to  $x$  and whose velocity is close to  $v$ .

Furthermore, with usual notations,  $E$  is the electric field (generated by electrons and ions themselves), while  $b$  is a constant external magnetic field. We assume that  $b = e_1$ , and consider  $(e_1, e_2, e_3)$  an orthonormal basis of  $\mathbb{R}^3$ .

It is interesting to notice that for a very particular class of (singular) data, namely Dirac functions in velocity, which we shall call *monokinetic* data,

$$f(t, x, v) = \rho(t, x)\delta_{v=u(t, x)}, \quad (1.3)$$

where  $\delta$  denotes the Dirac delta function, we get that  $f$  is a solution in the sense of distributions to (1.2) (with some relevant electric field) if and only if  $(\rho, u)$  satisfies the following hydrodynamic system, which corresponds to the *pressureless* Euler-Poisson system:

$$\begin{cases} \partial_t \rho + \nabla_x \cdot (\rho u) = 0, \\ \partial_t u + u \cdot \nabla_x u = E + u \wedge b, \quad u = (u_1, u_2, u_3), \\ E = -\nabla_x \phi, \\ -\Delta_x \phi + e^\phi = \rho. \end{cases} \quad (1.4)$$

From the physical point of view, this corresponds to the *cold ions* assumption, that corresponds to the situation where ions have zero (kinetic) temperature: any function of the form (1.3) indeed satisfies

$$\int f(t, x, v)|v - u(t, x)|^2 dv = 0. \quad (1.5)$$

We also mention that a standard model in plasma physics is obtained after linearizing the Maxwell-Boltzmann law, which yields from (1.2) the following system:

$$\begin{cases} \partial_t f + v \cdot \nabla_x f + E \cdot \nabla_v f = 0, \\ E = -\nabla_x \phi, \\ -\Delta_x \phi + \phi = \int_{\mathbb{R}^d} f dv - 1, \\ f|_{t=0} = f_0. \end{cases} \quad (1.6)$$

This linearization produces some considerable simplifications in the analysis.

*1.2. The long wave scaling.* In some recent independent works, Lannes, Linares and Saut [29] in the first hand, and Guo and Pu [22, 38] in the other hand, have rigorously studied the *long wave limit* of the pressureless Euler-Poisson system. Precisely, one looks for solutions to (1.4) under the form:

$$\begin{cases} \rho = 1 + \varepsilon \rho_1(\varepsilon^{1/2}(x_1 - t), \varepsilon^{1/2}x_2, \varepsilon^{1/2}x_3, \varepsilon^{3/2}t) + \varepsilon^2 \rho_2 + \dots, \\ \phi = \varepsilon \phi_1(\varepsilon^{1/2}(x_1 - t), \varepsilon^{1/2}x_2, \varepsilon^{1/2}x_3, \varepsilon^{3/2}t) + \varepsilon^2 \phi_2 + \dots, \\ u_1 = \varepsilon u_1^{(1)}(\varepsilon^{1/2}(x_1 - t), \varepsilon^{1/2}x_2, \varepsilon^{1/2}x_3, \varepsilon^{3/2}t) + \varepsilon^2 u_1^{(2)} + \dots, \\ u_2 = \varepsilon^{3/2} u_2^{(1)}(\varepsilon^{1/2}(x_1 - t), \varepsilon^{1/2}x_2, \varepsilon^{1/2}x_3, \varepsilon^{3/2}t) + \varepsilon^2 u_2^{(2)} + \dots, \\ u_3 = \varepsilon^{3/2} u_3^{(1)}(\varepsilon^{1/2}(x_1 - t), \varepsilon^{1/2}x_2, \varepsilon^{1/2}x_3, \varepsilon^{3/2}t) + \varepsilon^2 u_3^{(2)} + \dots \end{cases} \quad (1.7)$$

This leads to the study of the behaviour, as the parameter  $\varepsilon$  goes to 0, of the solutions to the rescaled system:

$$\begin{cases} \varepsilon \partial_t \rho - \partial_{x_1} \rho + \nabla_x \cdot ((1 + \varepsilon \rho)u) = 0, \\ \varepsilon \partial_t u - \partial_{x_1} u + \varepsilon u \cdot \nabla_x u = E + \varepsilon^{-1/2} u \wedge b, & u = (u_1, u_2, u_3), \\ E = -\nabla_x \phi, \\ -\varepsilon^2 \Delta_x \phi + e^{\varepsilon \phi} = 1 + \varepsilon \rho. \end{cases} \quad (1.8)$$

In the limit, one expects to obtain some (non-linear) dispersive equations, as explained in the papers by Zakharov-Kuznetsov [45] and Laedke-Spatschek [28]. Indeed, in  $1D$ , both [29] and [22] establish the convergence to the Korteweg-de Vries (KdV) equation, which we recall below:

$$\partial_t \phi_1 + \phi_1 \partial_x \phi_1 + \partial_{xxx}^3 \phi_1 = 0. \quad (1.9)$$

In  $2D$  and  $3D$ , which corresponds to the setting of (1.8), [29] and [38] give the derivation of a higher dimensional generalization of the KdV equation, which is referred to as the Zakharov-Kuznetsov (in short ZK) equation:

$$\partial_t \phi_1 + \phi_1 \partial_x \phi_1 + \partial_{x_1} \Delta \phi_1 = 0. \quad (1.10)$$

With another kind of anisotropic (in space) scaling, Pu has also derived in  $2D$  the Kadomstev-Petviashvili II (in short KP-II) equation [38]:

$$\partial_{x_1} (\partial_t \phi_1 + \phi_1 \partial_{x_1} \phi_1 + \partial_{x_1 x_1 x_1}^3 \phi_1) + \partial_{x_2 x_2}^2 \phi_1 = 0. \quad (1.11)$$

Over the past years, there have been many mathematical studies of long wave limits towards KdV (and higher dimensional generalizations). Let us cite some works (this list is not meant to be exhaustive) concerning the following PDEs:

- *Nonlinear Schrödinger (including Gross-Pitaevskii) equations*: Chiron and Rousset [14], and Béthuel, Gravejat, Saut and Smets [6,7] (with different methods),
- *General Hyperbolic systems*: Ben-Youssef and Colin [4], Ben-Youssef and Lannes [5],
- *Water Waves*: Craig [16], Schneider and Wayne [43], Bona, Colin and Lannes [8], Alvarez-Samaniego and Lannes [2], Duchene [17,18],

and many others.

The fact that there exist very singular solutions to the Vlasov-Poisson system (1.2) which precisely yield the pressureless Euler-Poisson system (1.4) suggests that it should also be possible to study (1.2) in a long wave regime. Following this idea, we would like to look for solutions of the form:

$$\begin{cases} f_\varepsilon(t, x, v) = \varepsilon^d \tilde{f}_\varepsilon(\varepsilon^{3/2} t, \varepsilon^{1/2}(x_1 - t), \varepsilon^{1/2} x_2, \varepsilon^{1/2} x_3, \varepsilon^{-1} v), \\ \phi_\varepsilon(x, v) = \varepsilon \tilde{\phi}_\varepsilon(\varepsilon^{3/2} t, \varepsilon^{1/2}(x_1 - t), \varepsilon^{1/2} x_2, \varepsilon^{1/2} x_3). \end{cases} \quad (1.12)$$

The normalization is chosen in order that the scaling of the two first moments  $\rho_\varepsilon := \int f_\varepsilon dv$  and  $u_\varepsilon := \frac{\int f_\varepsilon v dv}{\int f_\varepsilon dv}$  matches with the Ansatz in (1.7).

At some point, we will also have to somehow impose that the function  $\tilde{f}_\varepsilon$  is “close” to a Dirac function, in order to reach the Euler-Poisson dynamics.

Therefore, we propose a long wave scaling for the Vlasov-Poisson equation (in dimension  $d = 3$ ), which we introduce now:

$$\begin{cases} \tilde{t} = \varepsilon^{3/2}t, & \tilde{x} = \varepsilon^{1/2}x, & \tilde{v} = \varepsilon^{-1}v, \\ \tilde{f}(\tilde{t}, \tilde{x}, \tilde{v}) = \varepsilon^{-3}f(t, x, v), \\ \tilde{\phi}(\tilde{t}, \tilde{x}) = \varepsilon^{-1}\phi(t, x), \\ \tilde{E}(\tilde{t}, \tilde{x}) = \varepsilon^{-3/2}E(t, x). \end{cases} \quad (1.13)$$

With a slight abuse of notation (we forget the tildes), we obtain the rescaled equations:

$$\begin{cases} \varepsilon^{3/2}\partial_t f + \varepsilon^{3/2}v \cdot \nabla_x f + \varepsilon^{-1}(\varepsilon^{3/2}E + \varepsilon v \wedge b) \cdot \nabla_v f = 0, \\ E = -\nabla_x \phi, \\ -\varepsilon^2 \Delta_x \phi + e^{\varepsilon \phi} = \int_{\mathbb{R}^3} f dv, \\ f|_{t=0} = f_0. \end{cases} \quad (1.14)$$

Finally, there only remains to perform the shift with respect to the first spatial variable:


$$\begin{cases} \bar{x}_1 = x_1 - t/\varepsilon, \\ \bar{f}(t, \bar{x}_1, x_2, x_3, v) = f(t, x, v), \\ \bar{\phi}(t, \bar{x}_1, x_2, x_3) = \phi(t, \bar{x}_1, x_2, x_3), \\ \bar{E}(t, \bar{x}_1, x_2, x_3) = E(t, \bar{x}_1, x_2, x_3). \end{cases} \quad (1.15)$$

With another abuse of notation, we end up with the rescaled Vlasov-Poisson system:

$$\begin{cases} \varepsilon \partial_t f_\varepsilon - \partial_{x_1} f_\varepsilon + \varepsilon v \cdot \nabla_x f_\varepsilon + \left( E_\varepsilon + \frac{v \wedge b}{\sqrt{\varepsilon}} \right) \cdot \nabla_v f_\varepsilon = 0, \\ E_\varepsilon = -\nabla_x \phi_\varepsilon, \\ -\varepsilon^2 \Delta_x \phi_\varepsilon + e^{\varepsilon \phi_\varepsilon} = \int_{\mathbb{R}^3} f_\varepsilon dv, \\ f_{\varepsilon, |t=0} = f_{\varepsilon, 0}. \end{cases} \quad (1.16)$$

To the best of our knowledge, although this scaling seems very natural, it has never been introduced yet, even in the physics literature. Our goal in this work is to study the behaviour, as  $\varepsilon \rightarrow 0$ , of solutions to (1.16), that also get close (in some sense that we shall precise later) at initial time to monokinetic data. According to the previous discussion, it is natural to expect to obtain the ZK equation in the limit.

The relations between the different systems are summarized in the following diagram:



Let us briefly comment on this picture. As already explained, the link between Vlasov-Poisson and Pressureless Euler-Poisson is given by monokinetic type of data. Unfortunately, these data are way too singular to fit in the known Cauchy theories (although some results for measure data are actually available in  $1D$ , we refer to the work of Zheng and Majda [46]). As a matter of fact, we are not even aware of any result proving rigorously the convergence to the pressureless Euler-Poisson system for data that would converge in some sense to monokinetic data; the stability estimates that would be needed are indeed missing.

As already said, the long wave limit from the Pressureless Euler-Poisson system has been performed in [29, 22, 38]. One important step is to build a solution in an interval of time which is independent of  $\varepsilon$ . In order to study the long wave limit of the Vlasov-Poisson equation, instead of trying to derive the Pressureless Euler-Poisson system, the idea is to perform *simultaneously* the cold ions and long wave limits. We shall start from global weak solutions to the Vlasov-Poisson equation, and therefore we will not have to face the difficulty of finding uniform lifespans.

To prove such a result, we shall rely on a classical energy method, namely the relative entropy method. The idea originates in the work of Yau [44] on the hydrodynamic limit of some Ginzburg-Landau equation. It was independently brought in kinetic theory by Golse in [10] (in the context of the incompressible Euler limit of the Boltzmann equation) and by Brenier in [13] (in the context of the quasineutral limit of the Vlasov-Poisson equation).

The basic principle of the relative entropy strategy consists in *modulating* some well-chosen functional that has to be conserved or dissipated by the physical system (for instance, the good choice is the entropy for the Boltzmann equation). The modulation is obtained in terms of the solution to the target equation. One has to ensure that this new functional allows to “measure” in a certain sense the distance between the solution to the original system and that to the target equation. Then one has to prove that the functional that has been constructed is a Lyapunov function for the system: this follows from exact computations and algebraic identities. The computations can be more or less lengthy and tedious. It can be also very technical to justify these.

The estimates which can be obtained have to be understood as *stability* estimates: for instance, the results proved in this work can be interpreted as the stability of monokinetic data, in the long wave regime, with a dynamics dictated by some the KdV or ZK equations. This also strongly suggests that for our long wave limit, there are stability phenomena (a la Penrose) at stake, exactly like for the case of the quasineutral limit (the effects of instabilities for the latter limit

are briefly discussed for instance in [21]); more precisely, two stream instabilities are bound to destabilize the system and make the long wave limit fail (but these are avoided when one considers monokinetic data). On the topic of instabilities for the Vlasov-Poisson system, we refer to [24, 23, 25, 30, 31, 37, 32, 33].

For the Vlasov equation, this method is remarkably well adapted to handle the cold ions limit, in other words the “convergence” to monokinetic data, as it will be clear later. This was observed for the study of the quasineutral limit of the Vlasov-Poisson system for electrons with fixed ions, as done by Brenier [13] (this was completed later by Masmoudi [36], see also [20]). More recently, in [26], we have studied the quasineutral limit of the Vlasov-Poisson equation for ions with small mass electrons (which corresponds to (1.2)). In that work, we have observed that this equation displays a  $L \log L$  structure that is reminiscent of that of the Boltzmann equation (we refer to the works of Saint-Raymond [40, 42], [41] for the incompressible Euler limit, see also the recent paper of Allemand [1]); this will also play a crucial role in this work.

It is worth noticing that the method provided in this paper can also be applied to study the KdV limit of the Euler-Poisson system, for data with only low uniform regularity. Indeed we can start from the global weak solutions built by Cordier and Peng [15] and use similar computations as in the present paper.

To conclude this introduction, let us mention that Haragus, Nicholls and Sattinger in [27] relied on the KdV approximation of the Euler-Poisson system to study (formally and numerically) the interaction of solitary waves. It would be very interesting to start an analogous program for the Vlasov-Poisson equation.

*1.3. Organization of the paper.* The paper is organized as follows: in Section 2, we provide the derivation of the KdV equation (see Theorem 3), starting from the 1D Vlasov-Poisson equation with a linearized Maxwell-Boltzmann law. The exposure of this relatively simple case will allow us to lay down the basic principles of the relative entropy method applied to the long wave limit. In addition to this pedagogical interest, the existence of global solutions to the KdV equation allows to give stability estimates which are valid for all times. In Section 3, we present the main result of this paper, which is the derivation of the ZK equation, in Theorem 6, starting from the 3D Vlasov-Poisson equation with the full Maxwell-Boltzmann law. The proof will be much more technical, in particular due to the fact that only a  $L \log L$  type of control is available for the electric potential (instead of a  $L^2$  bound which can be obtained with a linearized Maxwell-Boltzmann law). We will need an unusually large number of correctors in the relative entropy. Finally, we give in two appendices some variants of our results (which can still be obtained with the relative entropy method): in particular we present another scaling for the 2D Vlasov-Poisson system which yields the KP-II equation in the long wave limit.

## 2. From the Vlasov-Poisson equation to the Korteweg-de Vries equation

In this section, we shall study the long wave limit of the 1D Vlasov-Poisson system with a linearized Maxwell-Boltzmann law, that is (here  $(x, v) \in \mathbb{T} \times \mathbb{R}$ ):

$$\begin{cases} \varepsilon \partial_t f_\varepsilon - \partial_x f_\varepsilon + \varepsilon v \partial_x f_\varepsilon + E_\varepsilon \partial_v f_\varepsilon = 0, \\ E_\varepsilon = -\partial_x \phi_\varepsilon, \\ -\varepsilon^2 \partial_{xx}^2 \phi_\varepsilon + \varepsilon \phi_\varepsilon = \int_{\mathbb{R}} f_\varepsilon dv - 1, \\ f_{\varepsilon, |t=0} = f_{\varepsilon, 0}. \end{cases} \quad (2.1)$$

*2.1. Preliminaries.* This system possesses an energy, which is conserved, at least formally:

$$\mathcal{E}_\varepsilon(t) := \frac{1}{2} \int f_\varepsilon |v|^2 dv dx + \frac{1}{2} \varepsilon \int |\partial_x \phi_\varepsilon|^2 dx + \frac{1}{2} \int |\phi_\varepsilon|^2 dx. \quad (2.2)$$

Using this energy, as well as the conservation of  $L_{x,v}^p$  norms that can be obtained using the hamiltonian structure of the Vlasov equation, one can prove, following the work of Arsenev [3], the following theorem, which states the existence of global weak solutions to (2.1):

**Theorem 1.** *Let  $\varepsilon > 0$ . Let  $f_{\varepsilon,0} \in L^1 \cap L^\infty(\mathbb{T} \times \mathbb{R})$  be a non-negative function such that the initial energy is bounded:*

$$\mathcal{E}_\varepsilon(0) := \frac{1}{2} \int f_{\varepsilon,0} |v|^2 dv dx + \frac{1}{2} \varepsilon \int |\partial_x \phi_{\varepsilon,0}|^2 dx + \frac{1}{2} \int |\phi_{\varepsilon,0}|^2 dx < +\infty, \quad (2.3)$$

where the initial electric potential  $\phi_{\varepsilon,0}$  is given by the elliptic equation:

$$-\varepsilon^2 \partial_{xx}^2 \phi_{\varepsilon,0} + \varepsilon \phi_{\varepsilon,0} = \int_{\mathbb{R}} f_{\varepsilon,0} dv - 1.$$

We also assume that:

$$\int f_{\varepsilon,0} dv dx = 1.$$

Then there exists a non-negative global weak solution  $f_\varepsilon \in L_t^\infty(L^1 \cap L^\infty(\mathbb{T} \times \mathbb{R}))$  to (2.1), such that the energy is non-increasing:

$$\forall t \geq t', \quad \mathcal{E}_\varepsilon(t) \leq \mathcal{E}_\varepsilon(t'), \quad (2.4)$$

and such that the following local conservation laws for  $(\rho_\varepsilon := \int f_\varepsilon dv, J_\varepsilon := \int f_\varepsilon v dv)$  are satisfied:

$$\partial_t \rho_\varepsilon - \frac{1}{\varepsilon} \partial_x \rho_\varepsilon + \partial_x J_\varepsilon = 0, \quad (2.5)$$

$$\partial_t J_\varepsilon - \frac{1}{\varepsilon} \partial_x J_\varepsilon + \partial_x \left( \int |v|^2 f_\varepsilon dv \right) = -\frac{1}{2} \partial_x (\phi_\varepsilon + 1)^2 + \frac{\varepsilon}{2} \partial_x |\partial_x \phi_\varepsilon|^2. \quad (2.6)$$

For the KdV equation, that we recall below,

$$\partial_t \phi_1 + \phi_1 \partial_x \phi_1 + \partial_{xxx}^3 \phi_1 = 0, \quad (2.7)$$

we have in hand a famous global existence result which was proved in the seminal paper of Bourgain [11]:

**Theorem 2.** *The KdV equation is globally well-posed in  $H^s(\mathbb{T})$  for  $s \geq 0$ .*

We will use this result only for large values of  $s$ .

*2.2. Formal derivation.* It is quite enlightening to perform a formal analysis in this simple one-dimensional case, to understand how the KdV equation arises. Formally, one directly considers monokinetic data:

$$f_\varepsilon(t, x, v) = \rho_\varepsilon(t, x) \delta_{v=u_\varepsilon(t, x)}. \quad (2.8)$$

Then, as already explained, we obtain the following pressureless Euler-Poisson equation in a long wave scaling:

$$\begin{cases} \partial_t \rho_\varepsilon - \frac{1}{\varepsilon} \partial_x \rho_\varepsilon + \partial_x (\rho_\varepsilon u_\varepsilon) = 0, \\ \partial_t u_\varepsilon - \frac{1}{\varepsilon} \partial_x u_\varepsilon + u_\varepsilon \partial_x u_\varepsilon = \frac{1}{\varepsilon} E_\varepsilon, \\ E = -\partial_x \phi_\varepsilon, \\ -\varepsilon^2 \partial_{xx}^2 \phi_\varepsilon + \varepsilon \phi_\varepsilon = \rho_\varepsilon - 1, \end{cases} \quad (2.9)$$

We look for an approximate solution satisfying the Ansatz:

$$\begin{cases} \rho_\varepsilon = 1 + \varepsilon \rho_1 + \mathcal{O}(\varepsilon^2), \\ \phi_\varepsilon = \phi_1 + \varepsilon \phi_2 + \mathcal{O}(\varepsilon^2), \\ u_\varepsilon = u_1 + \varepsilon u_2 + \mathcal{O}(\varepsilon^2). \end{cases} \quad (2.10)$$

Plugging this Ansatz in the equations, and matching the different powers of  $\varepsilon$ , we obtain a cascade of equations.

– Conservation of charge equation:

$$\mathcal{O}(1) : \quad -\partial_x \rho_1 + \partial_x u_1 = 0. \quad (2.11)$$

$$\mathcal{O}(\varepsilon) : \quad \partial_t \rho_1 - \partial_x \rho_2 + \partial_x (\rho_1 u_1) + \partial_x u_2 = 0. \quad (2.12)$$

– Momentum equation:

$$\mathcal{O}(\varepsilon^{-1}) : \quad -\partial_x u_1 + \partial_x \phi_1 = 0, \quad (2.13)$$

$$\mathcal{O}(1) : \quad \partial_t u_1 - \partial_x u_2 + u_1 \partial_x u_1 + \partial_x \phi_2 = 0. \quad (2.14)$$

– Poisson equation

$$\mathcal{O}(\varepsilon) : \quad \phi_1 = \rho_1, \quad (2.15)$$

$$\mathcal{O}(\varepsilon^2) : \quad -\partial_{xx}^2 \phi_1 + \phi_2 = \rho_2. \quad (2.16)$$



We clearly get from (2.11) and (2.13):

$$\phi_1 = \rho_1 = u_1 \quad (2.17)$$

and in the other hand from (2.14) and (2.16):

$$\begin{aligned} \partial_x(\phi_2 - u_2) &= \partial_x(\phi_2 - \rho_2) + \partial_x(\rho_2 - u_2) \\ &= \partial_{xxx}^2 \phi_1 + \partial_t u_1 + 2u_1 \partial_x u_1. \end{aligned} \quad (2.18)$$

Hence, from (2.12) and (2.18), we conclude that  $\phi_1$  satisfies the KdV equation:

$$2\partial_t \phi_1 + 3\phi_1 \partial_x \phi_1 + \partial_{xxx}^3 \phi_1 = 0. \quad (2.19)$$

*2.3. Rigorous derivation.* We now state the theorem which rigorously proves the convergence to KdV.

**Theorem 3.** *Let  $(f_{\varepsilon,0})_{\varepsilon \in (0,1)}$  be a family of non-negative initial data satisfying the assumptions of Theorem 1 and such that there exists  $C > 0$  with:*

$$\mathcal{E}_\varepsilon(0) \leq C, \quad \forall \varepsilon \in (0,1). \quad (2.20)$$

*We denote by  $(f_\varepsilon)_{\varepsilon \in (0,1)}$  a family of non-negative global weak solutions to (2.1) given by Theorem 1. Let  $\mathcal{H}_\varepsilon$  be the relative entropy defined by the functional:*

$$\begin{aligned} \mathcal{H}_\varepsilon(t) &:= \frac{1}{2} \int f_\varepsilon |v - u_1 - \varepsilon u_2|^2 dv dx + \frac{1}{2} \varepsilon \int |\partial_x \phi_\varepsilon - \partial_x \phi_1 - \varepsilon \partial_x \phi_2|^2 dx \\ &\quad + \frac{1}{2} \int (\phi_\varepsilon - \phi_1 - \varepsilon \phi_2)^2 dx, \end{aligned} \quad (2.21)$$

*where  $(u_1, \phi_1, u_2, \phi_2) \in [C([0, +\infty[, H^{s+2}(\mathbb{T}))]^4$  (with  $s > 3/2$ ) satisfy the following system:*

$$\begin{cases} 2\partial_t \phi_1 + 3\phi_1 \partial_x \phi_1 + \partial_{xxx}^3 \phi_1 = 0, \\ u_1 = \phi_1, \\ \partial_x(u_2 - \phi_2) = \partial_t \phi_1 + \phi_1 \partial_x \phi_1. \end{cases} \quad (2.22)$$

*Then there exist  $C_1, C_2 > 0$ , such that for all  $\varepsilon \in (0,1)$ ,*

$$\forall t \geq 0, \quad \mathcal{H}_\varepsilon(t) \leq \mathcal{H}_\varepsilon(0) + \int_0^t (C_1 \mathcal{H}_\varepsilon(s) + C_2 \sqrt{\varepsilon}) ds. \quad (2.23)$$

*Assuming in addition that there exists  $C_3$  such that for any  $\varepsilon \in (0,1)$ :*

$$\mathcal{H}_\varepsilon(0) \leq C_3 \sqrt{\varepsilon}, \quad (2.24)$$

*then we obtain for any  $\varepsilon \in (0,1)$ :*

$$\forall t \geq 0, \quad \mathcal{H}_\varepsilon(t) \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}. \quad (2.25)$$

In order to get functions satisfying (2.22), we can proceed as follows:

– The existence of  $\phi_1$  is ensured by Theorem 2.

- We accordingly set  $\phi_1 = u_1$ .
- The functions  $\phi_2$  and  $u_2$  can be seen as correctors. The last equation of (2.22) allows to define them. Only the value of the function  $u_2 - \phi_2$  is important.

*Remark 1.* The estimate (2.25) clearly implies that

$$\frac{1}{2} \int f_\varepsilon |v - u_1 - \varepsilon u_2|^2 dv dx \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}, \quad (2.26)$$

and

$$\frac{1}{2} \int (\phi_\varepsilon - \phi_1 - \varepsilon \phi_2)^2 dx \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}. \quad (2.27)$$

From (2.26), denoting by  $f$  a weak limit of  $f_\varepsilon$ , we deduce that necessarily

$$\int f |v - u_1|^2 dv dx = 0, \quad (2.28)$$

which means that the limit temperature is zero (cold ions limit).

*Remark 2.* The estimate (2.25) is valid for all times, but it is useful for times of order  $o(|\log \varepsilon|)$  (in other words, logarithmically growing times). This is slightly better than the times of validity obtained in the long wave limit of Euler-Poisson in [29], which are (translated in our framework) of order  $O(1)$ .

From this theorem, we can deduce the following corollary:

**Corollary 1.** *Making the same assumptions as in the previous theorem, we obtain the weak convergences:*

$$\begin{cases} \rho_\varepsilon \rightharpoonup_{\varepsilon \rightarrow 0} 1 \text{ in } L_t^\infty \mathcal{M}^1 \text{ weak-}^*, \\ J_\varepsilon \rightharpoonup_{\varepsilon \rightarrow 0} \phi_1 \text{ in } L_t^\infty \mathcal{M}^1 \text{ weak-}^*, \\ \phi_\varepsilon \rightharpoonup_{\varepsilon \rightarrow 0} \phi_1 \text{ in } L_t^\infty \mathcal{M}^1 \text{ weak-}^*. \end{cases} \quad (2.29)$$

*Proof (Proof of Corollary 1).*

By conservation of the  $L^1$  norm, we deduce that

$$\|\rho_\varepsilon\|_{L^1} \leq C.$$

Using this bound and the one coming from the energy inequality:

$$\int f_\varepsilon |v|^2 dv dx \leq C,$$

and by non-negativity of  $f_\varepsilon$ , it is standard to deduce a  $L^1$  control on  $J_\varepsilon$ :

$$\|\rho_\varepsilon\|_{L^1} \leq C.$$

Therefore, there exist two non-negative measures  $\rho, J$  such that up to some extraction, we have:

$$\begin{cases} \rho_\varepsilon \rightharpoonup \rho, \\ J_\varepsilon \rightharpoonup J, \end{cases} \quad (2.30)$$

in the vague sense of measures. We have to show that  $\rho = 1$  and  $J = \phi_1$ .

We pass to the limit  $\varepsilon \rightarrow 0$  (in the sense of distributions) in the Poisson equation:

$$-\varepsilon^2 \partial_{xx}^2 \phi_\varepsilon + \varepsilon \phi_\varepsilon = \rho_\varepsilon - 1,$$

using the uniform  $L^2$  bound on  $\sqrt{\varepsilon} \partial_x \phi_\varepsilon$  and  $\phi_\varepsilon$  (coming from the energy inequality), we deduce that we necessarily have  $\rho = 1$ , which proves the first claim.

In the other hand, using the Cauchy-Schwarz inequality, we have

$$\frac{|J_\varepsilon - \rho_\varepsilon(u_1 + \varepsilon u_2)|^2}{\rho_\varepsilon} = \frac{\left(\int f_\varepsilon(v - u_1 - \varepsilon u_2) dv\right)^2}{\int f_\varepsilon dv} \leq \int f_\varepsilon |v - u_1 - \varepsilon u_2|^2 dv, \quad (2.31)$$

The functional  $(\rho, J) \rightarrow \int \frac{|J - \rho(u_1 + \varepsilon u_2)|^2}{\rho} dx$  is convex and lower semi-continuous with respect to the weak convergence of measures (see [13]). As a consequence, the weak convergences in the vague sense of measures  $\rho_\varepsilon \rightarrow 1$  and  $J_\varepsilon \rightarrow J$  lead to:

$$\int |J - u_1|^2 dx \leq \liminf_{\varepsilon \rightarrow 0} \int \frac{|J_\varepsilon - \rho_\varepsilon u|^2}{\rho_\varepsilon} dx. \quad (2.32)$$

By (2.25), we deduce that  $J = u_1 (= \phi_1)$ .

To conclude, the uniqueness of the limit allows us to say that the weak convergences actually hold without any extraction.

*Remark 3.* We can actually state strong convergence results. Indeed, in view of the preceding proof, it is clear that (2.26) implies the “strong” convergence:

$$\left\| \frac{J_\varepsilon - \rho_\varepsilon(u_1 + \varepsilon u_2)}{\sqrt{\rho_\varepsilon}} \right\|_{L^2}^2 \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}. \quad (2.33)$$

In the other hand, the control (2.27) means that  $\phi_\varepsilon$  converges strongly in  $L^2$  to  $\phi_1$ , as  $\varepsilon$  goes to 0.

*2.4. Proof of Theorem 3.* Relying on the fact the energy is non-increasing (and that  $\mathcal{H}_\varepsilon$  is built as a modulation of the energy), we have for all  $t \geq 0$  and all  $\varepsilon \in (0, 1)$ :

$$\mathcal{H}_\varepsilon(t) = \mathcal{E}_\varepsilon(t) + (\mathcal{H}_\varepsilon(t) - \mathcal{E}_\varepsilon(t)) \leq \mathcal{E}_\varepsilon(0) + (\mathcal{H}_\varepsilon(t) - \mathcal{E}_\varepsilon(t)),$$

which yields:

$$\begin{aligned} \mathcal{H}_\varepsilon(t) &\leq \mathcal{H}_\varepsilon(0) + \int_0^t \int \partial_t \left[ f_\varepsilon \left( \frac{1}{2} |u_1 + \varepsilon u_2|^2 - v(u_1 + \varepsilon u_2) \right) \right] dv dx \\ &\quad + \varepsilon \int \partial_t \left[ \frac{1}{2} |\partial_x \phi_1 + \varepsilon \partial_x \phi_2|^2 - \partial_x \phi_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) \right] dx \\ &\quad + \int \partial_t \left[ \frac{1}{2} |\phi_1 + \varepsilon \phi_2|^2 - \phi_\varepsilon (\phi_1 + \varepsilon \phi_2) \right] dx ds \\ &:= \mathcal{H}_\varepsilon(0) + \int_0^t (I_1 + I_2 + I_3) ds. \end{aligned}$$

We are now going to study  $I_1$ ,  $I_2$  and  $I_3$ . The computations can be justified using only the local conservation laws (2.5) and (2.6).

**Study of  $I_1$ .**

Using the fact that  $f_\varepsilon$  satisfies the Vlasov-Poisson equation, we obtain the identity:

$$\begin{aligned} & \int \partial_t f_\varepsilon \left( \frac{1}{2} |u_1 + \varepsilon u_2|^2 - v(u_1 + \varepsilon u_2) \right) dv dx \\ &= \int f_\varepsilon (u_1 + \varepsilon u_2 - v) \left[ -\frac{1}{\varepsilon} \partial_x u_1 + v \partial_x u_1 - \partial_x u_2 + \varepsilon v \partial_x u_2 \right] dv dx \\ & \quad - \int \frac{1}{\varepsilon} \rho_\varepsilon E_\varepsilon u_1 dx - \int \rho_\varepsilon E_\varepsilon u_2 dx. \end{aligned}$$

In order to obtain an hydrodynamic equation inside [...], we write:

$$\begin{aligned} \int f_\varepsilon (u_1 + \varepsilon u_2 - v) (v \partial_x u_1) &= \int f_\varepsilon (u_1 + \varepsilon u_2 - v) (u_1 \partial_x u_1) \\ & \quad - \int f_\varepsilon |u_1 + \varepsilon u_2 - v|^2 \partial_x u_1 + \int \varepsilon f_\varepsilon (u_1 + \varepsilon u_2 - v) u_2 \partial_x u_1. \end{aligned}$$

After deriving with respect to time the term  $\frac{1}{2} |u_1 + \varepsilon u_2|^2 - v(u_1 + \varepsilon u_2)$ , we get the following contribution in  $I_1$ :

$$\int f_\varepsilon (u_1 + \varepsilon u_2 - v) \left[ \partial_t u_1 + \varepsilon \partial_t u_2 \right] dv dx.$$

We now focus on the terms of order  $\mathcal{O}(1/\varepsilon)$ , for which we can write

$$\begin{aligned} & - \int f_\varepsilon (u_1 + \varepsilon u_2 - v) \frac{1}{\varepsilon} \partial_x u_1 dv dx \\ & \quad = \int f_\varepsilon (u_1 + \varepsilon u_2 - v) \frac{1}{\varepsilon} (-\partial_x u_1 + \partial_x \phi_1) dv dx \\ & \quad \quad - \int \frac{1}{\varepsilon} \rho_\varepsilon u_1 \partial_x \phi_1 dx - \int \rho_\varepsilon u_2 \partial_x \phi_1 dx + \int \frac{1}{\varepsilon} J_\varepsilon \partial_x \phi_1 dx, \end{aligned}$$

and

$$\begin{aligned} & - \int \frac{1}{\varepsilon} \rho_\varepsilon E_\varepsilon u_1 dx - \int \frac{1}{\varepsilon} \rho_\varepsilon u_1 \partial_x \phi_1 dx - \int \rho_\varepsilon u_1 \partial_x \phi_2 dx \\ & \quad = -\frac{1}{\varepsilon} \int \rho_\varepsilon u_1 (\partial_x \phi_1 + \varepsilon \partial_x \phi_2 - \partial_x \phi) dx. \end{aligned}$$

In the other hand, we observe that we can write:

$$\begin{aligned} & \int f_\varepsilon (u_1 + \varepsilon u_2 - v) \left[ \partial_t u_1 + u_1 \partial_x u_1 - \partial_x u_2 \right] dv dx \\ & \quad = \int f_\varepsilon (u_1 + \varepsilon u_2 - v) \left[ \partial_t u_1 + u_1 \partial_x u_1 - \partial_x u_2 + \partial_x \phi_2 \right] dv dx \\ & \quad \quad + \int J_\varepsilon \partial_x \phi_2 dx - \int \rho_\varepsilon u_1 \partial_x \phi_2 dx - \varepsilon \int \rho_\varepsilon u_2 \partial_x \phi_2 dx. \end{aligned}$$

Using the  $L^1$  uniform bounds for  $\rho_\varepsilon$  and  $J_\varepsilon$ , as well as the various Lipschitz bounds on  $(u_1, u_2)$ , it is clear that

$$\left| \varepsilon \int \rho_\varepsilon u_2 \partial_x \phi_2 dx \right| + \left| \varepsilon \int f_\varepsilon(u_1 + \varepsilon u_2 - v) \partial_t u_2 dv dx \right| \leq C\varepsilon.$$

In the end, we have:

$$\begin{aligned} I_1 &= \frac{1}{\varepsilon} \int f_\varepsilon(u_1 + \varepsilon u_2 - v) \left[ -\partial_x u_1 + \partial_x \phi_1 \right] dv dx \\ &\quad + \int f_\varepsilon(u_1 + \varepsilon u_2 - v) \left[ \partial_t u_1 + v \partial_x u_1 - \partial_x u_2 + \partial_x \phi_2 \right] dv dx \\ &\quad + \int J_\varepsilon \left( \frac{1}{\varepsilon} \partial_x \phi_1 + \partial_x \phi_2 \right) dx \\ &\quad - \frac{1}{\varepsilon} \int \rho_\varepsilon u_1 (\partial_x \phi_1 + \varepsilon \partial_x \phi_2 - \partial_x \phi) dx \\ &\quad - \frac{1}{\varepsilon} \int \rho_\varepsilon u_2 \partial_x (\phi_1 + \varepsilon \phi_2) dx - \int \rho_\varepsilon E_\varepsilon u_2 dx \\ &\quad - \int f_\varepsilon |u_1 + \varepsilon u_2 - v|^2 \partial_x u_1 + \mathcal{O}(\varepsilon), \end{aligned} \tag{2.34}$$

where  $\mathcal{O}(\varepsilon)$  is a notation for all the terms which can be bounded by  $C\varepsilon$ , with  $C > 0$  independent of  $\varepsilon$ .

#### Study of $I_2$ .

We have:

$$\begin{aligned} I_2 &= \varepsilon \int \partial_t (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) (-\partial_x \phi_\varepsilon + \partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx \\ &\quad - \varepsilon \int \partial_t \partial_x \phi_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx. \end{aligned}$$

We get the easy bound (using  $|ab| \leq \frac{1}{2}(a^2 + b^2)$ ):

$$\begin{aligned} &\varepsilon \left| \int (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) (-\partial_x \phi_\varepsilon + \partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx \right| \\ &\leq C \left[ \varepsilon \int (\partial_t (\partial_x \phi_1 + \varepsilon \partial_x \phi_2))^2 dx + \varepsilon \int (-\partial_x \phi_\varepsilon + \partial_x \phi_1 + \varepsilon \partial_x \phi_2)^2 dx \right] \\ &\leq \mathcal{O}(\varepsilon) + C\mathcal{H}_\varepsilon(t). \end{aligned}$$

#### Study of $I_3$ .

We get:

$$\begin{aligned} I_3 &= \int \partial_t (\phi_1 + \varepsilon \phi_2) (-\phi_\varepsilon + \phi_1 + \varepsilon \phi_2) dx \\ &\quad - \int \partial_t \phi_\varepsilon (\phi_1 + \varepsilon \phi_2) dx := I_3^1 + I_3^2. \end{aligned}$$

Let us start with  $I_3^1$ , that we can rewrite as:

$$I_3^1 = \int (\phi_1 - \phi_\varepsilon) \partial_t \phi_1 + \varepsilon \int \partial_t \phi_2 (-\phi_\varepsilon + \phi_1 + \varepsilon \phi_2) dx + \varepsilon \int \partial_t \phi_1 \phi_2 dx.$$

If we differentiate with respect to time the Poisson equation, we obtain:

$$-\partial_t \phi_\varepsilon = -\varepsilon \partial_{xx}^2 \partial_t \phi_\varepsilon - \frac{1}{\varepsilon} \partial_t \rho_\varepsilon.$$

Then, using the local conservation of charge (in shifted variables) that we recall below,

$$-\frac{1}{\varepsilon} \partial_t \rho_\varepsilon = -\frac{1}{\varepsilon^2} \partial_x \rho_\varepsilon + \frac{1}{\varepsilon} \partial_x J_\varepsilon,$$

we obtain that:

$$\begin{aligned} I_3^2 &= -\frac{1}{\varepsilon^2} \int \partial_x \rho_\varepsilon (\phi_1 + \varepsilon \phi_2) dx \\ &\quad + \frac{1}{\varepsilon} \int \partial_x J_\varepsilon \phi_1 dx + \int \partial_x J_\varepsilon \phi_2 dx \\ &\quad - \varepsilon \int \partial_t \partial_{xx}^2 \phi_\varepsilon (\phi_1 + \varepsilon \phi_2). \end{aligned}$$

Of course, by integration by parts, we can rewrite the last three terms as follows:

$$\begin{aligned} &\frac{1}{\varepsilon} \int \partial_x J_\varepsilon \phi_1 dx + \int \partial_x J_\varepsilon \phi_2 dx + \varepsilon \int \partial_t \partial_{xx}^2 \phi_\varepsilon (\phi_1 + \varepsilon \phi_2) \\ &= -\frac{1}{\varepsilon} \int J_\varepsilon \partial_x \phi_1 dx - \int J_\varepsilon \partial_x \phi_2 dx + \varepsilon \int \partial_t \partial_x \phi_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2), \end{aligned}$$

which get simplified using some terms coming from the computations of  $I_1$  and  $I_2$ .

### Study of the remaining significant terms.

We now have to study the following terms (coming from the computation of  $I_1 + I_2 + I_3$ ):

$$\begin{aligned} K_1 &:= - \int \rho_\varepsilon u_2 \partial_x (\phi_1 + \varepsilon \phi_2) dx, & K_2 &:= - \int \rho_\varepsilon E_\varepsilon u_2 dx, \\ K_3 &:= -\frac{1}{\varepsilon} \int \rho_\varepsilon u_1 (\partial_x \phi_1 + \varepsilon \partial_x \phi_2 - \partial_x \phi_\varepsilon) dx, \\ K_4 &:= -\frac{1}{\varepsilon^2} \int \partial_x \rho_\varepsilon (\phi_1 + \varepsilon \phi_2) dx, \\ K_5 &:= \int (\phi_1 - \phi_\varepsilon) \partial_t \phi_1 dx. \end{aligned}$$

We start with  $K_1$  and  $K_2$ . Using the Poisson equation, we have:

$$\begin{aligned} K_1 &= - \int u_2 \partial_x (\phi_1 + \varepsilon \phi_2) dx - \varepsilon \int \phi_\varepsilon u_2 \partial_x (\phi_1 + \varepsilon \phi_2) dx \\ &\quad + \varepsilon^2 \int \partial_{xx}^2 \phi_\varepsilon u_2 \partial_x (\phi_1 + \varepsilon \phi_2) dx \\ &= \int \partial_x u_2 \phi_1 dx - \varepsilon \int \phi_\varepsilon u_2 \partial_x \phi_1 dx + \varepsilon^2 \int \partial_{xx}^2 \phi_\varepsilon u_2 \partial_x (\phi_1 + \varepsilon \phi_2) dx \\ &\quad + \mathcal{O}(\varepsilon). \end{aligned}$$

We have (after integration by parts),

$$\varepsilon^2 \left| \int \partial_{xx}^2 \phi_\varepsilon u_2 \partial_x (\phi_1 + \varepsilon \phi_2) dx \right| \leq C\varepsilon.$$

Similarly, we compute

$$\begin{aligned} K_2 &= \int u_2 \partial_x \phi_\varepsilon dx + \varepsilon \int \phi_\varepsilon u_2 \partial_x \phi_\varepsilon dx - \varepsilon^2 \int \partial_{xx}^2 \phi_\varepsilon u_2 \partial_x \phi_\varepsilon dx \\ &= - \int \partial_x u_2 \phi_\varepsilon dx - \varepsilon \int \frac{\phi_\varepsilon^2}{2} \partial_x u_2 dx + \varepsilon^2 \int \frac{(\partial_x \phi_\varepsilon)^2}{2} \partial_x u_2 dx. \end{aligned}$$

As a result, one gets

$$K_1 + K_2 = \int (\phi_1 - \phi_\varepsilon) \partial_x u_2 dx + \mathcal{O}(\varepsilon).$$

Concerning  $K_3$ , we have (once again using the Poisson equation):

$$\begin{aligned} K_3 &= -\frac{1}{\varepsilon} \int u_1 (\partial_x \phi_1 + \varepsilon \partial_x \phi_2 - \partial_x \phi_\varepsilon) dx \\ &\quad - \int \phi_\varepsilon u_1 (\partial_x \phi_1 + \varepsilon \partial_x \phi_2 - \partial_x \phi_\varepsilon) dx \\ &\quad + \int \varepsilon \partial_{xx}^2 \phi_\varepsilon u_1 (\partial_x \phi_1 + \varepsilon \partial_x \phi_2 - \partial_x \phi_\varepsilon) dx \\ &:= K_3^1 + K_3^2 + K_3^3. \end{aligned}$$

For  $K_3^1$ , we have:

$$K_3^1 = -\frac{1}{\varepsilon} \int u_1 \partial_x \phi_1 dx - \int u_1 \partial_x \phi_2 dx + \frac{1}{\varepsilon} \int u_1 \partial_x \phi_\varepsilon dx.$$

Concerning  $K_3^2$ , we write:

$$K_3^2 = - \int \phi_\varepsilon u_1 \partial_x u_1 dx + \int \phi_\varepsilon \partial_x \phi_\varepsilon u_1 dx + \varepsilon \int \phi_\varepsilon u_1 \partial_x \phi_2 dx$$

and

$$- \int \phi_\varepsilon u_1 \partial_x u_1 dx = \int (\phi_1 - \phi_\varepsilon) u_1 \partial_x u_1 dx - \int \phi_1 u_1 \partial_x u_1 dx.$$

The contribution coming from  $\int \phi_\varepsilon \partial_x \phi_\varepsilon u_1 dx$  could be harmful (a priori it is of order  $\mathcal{O}(1/\varepsilon)$ ), but we can rely on the following identities in order to make the relative entropy appear:

$$\begin{aligned}
& \int \phi_\varepsilon \partial_x \phi_\varepsilon u_1 dx = -\frac{1}{2} \int \phi_\varepsilon^2 \partial_x u_1 dx \\
& = -\frac{1}{2} \int (\phi_\varepsilon - \phi_1 - \varepsilon \phi_2)^2 \partial_x u_1 dx - \int \phi_\varepsilon \phi_1 \partial_x u_1 dx + \varepsilon \int \phi_\varepsilon \phi_2 \partial_x u_1 dx \\
& \quad + \frac{1}{2} \int (\phi_1 + \varepsilon \phi_2)^2 \partial_x u_1 dx \\
& = -\frac{1}{2} \int (\phi_\varepsilon - \phi_1 - \varepsilon \phi_2)^2 \partial_x u_1 dx + \int (\phi_1 - \phi_\varepsilon) \phi_1 \partial_x u_1 dx + \varepsilon \int \phi_\varepsilon \phi_2 \partial_x u_1 dx \\
& \quad - \frac{1}{2} \int \phi_1^2 \partial_x u_1 dx + \varepsilon \int \left( \phi_1 \phi_2 \partial_x u_1 + \frac{1}{2} \varepsilon \phi_2^2 \partial_x u_1 \right) dx.
\end{aligned}$$

We note that  $\varepsilon \int \phi_\varepsilon \phi_2 \partial_x u_1 dx$  is of order  $\varepsilon$ , using the  $L^2$  uniform bound on  $\phi_\varepsilon$  granted by the energy.

Finally, for  $K_3^3$ , we have

$$K_3^3 = - \int \partial_{xx}^2 \phi_\varepsilon u_1 \partial_x \phi_\varepsilon dx + \varepsilon \int \partial_{xx}^2 \phi_\varepsilon u_1 (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx.$$

As before, the most significant term can be rewritten as follows:

$$\begin{aligned}
& - \int \partial_{xx}^2 \phi_\varepsilon u_1 \partial_x \phi_\varepsilon dx = \varepsilon \int \frac{(\partial_x \phi_\varepsilon)^2}{2} \partial_x u_1 dx \\
& = \frac{1}{2} \int \varepsilon (\partial_x \phi_\varepsilon - \partial_x \phi_1 - \varepsilon \partial_x \phi_2)^2 \partial_x u_1 + \varepsilon \int \partial_x \phi_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) \partial_x u_1 dx \\
& \quad - \frac{1}{2} \int \varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2)^2 \partial_x u_1 dx.
\end{aligned}$$

We have the bound:

$$\varepsilon \left| \int \partial_x \phi_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) \partial_x u_1 dx \right| \leq C \sqrt{\varepsilon}.$$

We treat  $K_4$  exactly as  $K_3$ .

$$\begin{aligned}
K_4 & = \frac{1}{\varepsilon^2} \int \rho_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx \\
& = \frac{1}{\varepsilon^2} \int (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx + \frac{1}{\varepsilon} \int \phi_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx \\
& \quad - \int \partial_{xx}^2 \phi_\varepsilon (\partial_x \phi_1 + \varepsilon \partial_x \phi_2) dx \\
& := K_4^1 + K_4^2 + K_4^3.
\end{aligned}$$

Clearly, we have  $K_4^1 = 0$  and

$$K_4^2 = \frac{1}{\varepsilon} \int \phi_\varepsilon \partial_x \phi_1 dx + \int \phi_\varepsilon \partial_x \phi_2 dx,$$



and since  $\int \phi_1 \partial_{xxx}^3 \phi_1 dx = 0$ , we have:

$$K_4^3 = \int (\phi_1 - \phi_\varepsilon) \partial_{xxx}^3 \phi_1 dx - \varepsilon \int \partial_{xx}^2 \phi_\varepsilon \partial_x \phi_2 dx.$$

By integration by parts, it is clear that the second term of  $K_4^3$  is of order  $\sqrt{\varepsilon}$ .

**Conclusion.**

Gathering all pieces together, we obtain that:

$$\begin{aligned} \mathcal{H}_\varepsilon(t) &\leq \mathcal{H}_\varepsilon(0) + \int_0^t \left[ \int f_\varepsilon(u_1 + \varepsilon u_2 - v) \frac{1}{\varepsilon} (-\partial_x u_1 + \partial_x \phi_1) dv dx \right. \\ &+ \int f_\varepsilon(u_1 + \varepsilon u_2 - v) \left[ \partial_t u_1 + u_1 \partial_x u_1 - \partial_x u_2 + \partial_x \phi_2 \right] dv dx \\ &+ \frac{1}{\varepsilon} \int (u_1 - \phi_1) \partial_x \phi_\varepsilon dx \\ &+ \int (\phi_1 - \phi_\varepsilon) (\partial_t \phi_1 + \phi_1 \partial_x u_1 + u_1 \partial_x u_1 + \partial_{xxx}^3 \phi_1 + \partial_x u_2 - \partial_x \phi_2) dx \\ &- \int (u_1 - \phi_1) \partial_x \phi_2 - \int \phi_1 u_1 \partial_x u_1 dx - \frac{1}{2} \int \phi_1^2 \partial_x u_1 \\ &- \int f_\varepsilon |v - u_1 - \varepsilon u_2|^2 \partial_x u_1 dv dx - \frac{1}{2} \varepsilon \int |\partial_x \phi_\varepsilon - \partial_x \phi_1 - \varepsilon \partial_x \phi_2|^2 \partial_x u_1 dx \\ &- \frac{1}{2} \int (\phi_\varepsilon - \phi_1 - \varepsilon \phi_2)^2 \partial_x u_1 dx \\ &\left. + \mathcal{O}(\sqrt{\varepsilon}) \right] ds. \end{aligned}$$

We impose the following cancellations (to kill all singular terms):

$$\begin{cases} u_1 - \phi_1 = 0, \\ \partial_t u_1 + u_1 \partial_x u_1 - \partial_x u_2 + \partial_x \phi_2 = 0, \\ \partial_t \phi_1 + \phi_1 \partial_x u_1 + u_1 \partial_x u_1 + \partial_{xxx}^3 \phi_1 + \partial_x u_2 - \partial_x \phi_2 = 0. \end{cases} \quad (2.35)$$

These are consequences of the identity (2.22). Using the Lipschitz bound on  $u_1$  we end up with:

$$\mathcal{H}_\varepsilon(t) \leq \mathcal{H}_\varepsilon(0) + \int_0^t (C_1 \mathcal{H}_\varepsilon(t) + C_2 \sqrt{\varepsilon}) ds, \quad (2.36)$$

which then yields the claimed Gronwall type bound (2.25). The proof is consequently complete.

### 3. From the Vlasov-Poisson equation to the Zakharov-Kuznetsov equation

We perform the analysis in  $3D$ , but this can also be done in  $2D$ , in an almost similar way.

We study the behaviour, as  $\varepsilon \rightarrow 0$ , of the solutions to the following equation (for  $(x, v) \in \mathbb{T}^3 \times \mathbb{R}^3$ ):

$$\begin{cases} \varepsilon \partial_t f_\varepsilon - \partial_{x_1} f_\varepsilon + \varepsilon v \cdot \nabla_x f_\varepsilon + \left( E_\varepsilon + \frac{v \wedge e_1}{\sqrt{\varepsilon}} \right) \cdot \nabla_v f_\varepsilon = 0, \\ E_\varepsilon = -\nabla_x \phi_\varepsilon, \\ -\varepsilon^2 \Delta_x \phi_\varepsilon + e^{\varepsilon \phi_\varepsilon} = \int_{\mathbb{R}^3} f_\varepsilon dv, \\ f_{\varepsilon, |t=0} = f_{\varepsilon, 0}. \end{cases} \quad (3.1)$$

*3.1. Preliminaries.* This system possesses an energy, which is conserved, at least formally:

$$\begin{aligned} \mathcal{E}_\varepsilon(t) &:= \frac{1}{2} \int f_\varepsilon |v|^2 dv dx \\ &+ \frac{1}{2} \varepsilon \int |\nabla_x \phi_\varepsilon|^2 dx + \frac{1}{\varepsilon^2} \int (e^{\varepsilon \phi_\varepsilon} (\varepsilon \phi_\varepsilon - 1) + 1) dx, \end{aligned} \quad (3.2)$$

Note that the third term of this energy has a  $L \log L$  structure:

$$\frac{1}{\varepsilon^2} \int (e^{\varepsilon \phi_\varepsilon} (\varepsilon \phi_\varepsilon - 1) + 1) dx = \frac{1}{\varepsilon^2} \int (e^{\varepsilon \phi_\varepsilon} \log(e^{\varepsilon \phi_\varepsilon} / e^0) - e^{\varepsilon \phi_\varepsilon} + e^0) dx.$$

We have the following global existence theorem, which can be adapted from the work of Bouchut [9]:

**Theorem 4.** *Let  $\varepsilon > 0$ . Let  $f_{\varepsilon, 0} \in L^1 \cap L^\infty(\mathbb{T}^3 \times \mathbb{R}^3)$  be a non-negative function such that:*

$$\begin{aligned} \mathcal{E}_\varepsilon(0) &:= \frac{1}{2} \int f_{\varepsilon, 0} |v|^2 dv dx \\ &+ \frac{1}{2} \varepsilon \int |\nabla_x \phi_{\varepsilon, 0}|^2 dx + \frac{1}{\varepsilon^2} \int (e^{\varepsilon \phi_{\varepsilon, 0}} (\varepsilon \phi_{\varepsilon, 0} - 1) + 1) dx < +\infty, \end{aligned} \quad (3.3)$$

where the initial electric potential  $\phi_{\varepsilon, 0}$  is given by the elliptic equation:

$$-\varepsilon^2 \Delta_x \phi_{\varepsilon, 0} + e^{\varepsilon \phi_{\varepsilon, 0}} = \int_{\mathbb{R}^3} f_{\varepsilon, 0} dv.$$

We also assume that:

$$\int f_{\varepsilon, 0} dv dx = 1.$$

Then there exists a non-negative global weak solution  $f_\varepsilon \in L_t^\infty(L^1 \cap L^\infty(\mathbb{T}^3 \times \mathbb{R}^3))$  to (3.1), such that:

$$\forall t \geq t', \quad \mathcal{E}_\varepsilon(t) \leq \mathcal{E}_\varepsilon(t'), \quad (3.4)$$

and such that the following conservation laws for  $(\rho_\varepsilon := \int f_\varepsilon dv, J_\varepsilon := \int f_\varepsilon v dv)$  are satisfied:

$$\partial_t \rho_\varepsilon - \frac{1}{\varepsilon} \partial_{x_1} \rho_\varepsilon + \nabla_x \cdot J_\varepsilon = 0, \quad (3.5)$$

$$\begin{aligned} \partial_t J_\varepsilon - \frac{1}{\varepsilon} \partial_{x_1} J_\varepsilon + \nabla_x : \left( \int v \otimes v f_\varepsilon dv \right) &= \frac{1}{\varepsilon^{3/2}} J_\varepsilon \wedge e_1 \\ &- \frac{1}{\varepsilon} \nabla_x (e^{\varepsilon \phi_\varepsilon}) + \varepsilon \nabla_x : (\nabla_x V_\varepsilon \otimes \nabla_x V_\varepsilon) - \frac{\varepsilon}{2} \nabla_x |\nabla_x V_\varepsilon|^2. \end{aligned} \quad (3.6)$$

Let us now turn to the Cauchy problem for the ZK equation, that we recall now:

$$\partial_t \phi_1 + \phi_1 \partial_x \phi_1 + \partial_{x_1} \Delta \phi_1 = 0. \quad (3.7)$$

The only result in  $3D$  we are aware of for this equation concerns the case of the whole space  $\mathbb{R}^3$  (for results in  $2D$ , we refer to [19, 34]): in [35], Linares and Saut proved that ZK is locally well-posed in  $H^s(\mathbb{R}^3)$ , for  $s > 9/8$ , and more recently in [39], Ribaud and Vento showed that it is well-posed for  $s > 1$ . Their proofs are based on dispersive effects and can not be directly applied to the case of the torus  $\mathbb{T}^3$ . Using standard methods, we can nevertheless prove the easy theorem:

**Theorem 5.** *The ZK equation is locally well-posed in  $H^s(\mathbb{T}^3)$  for  $s > 5/2$ .*

This will be sufficient for our needs.

*3.2. Rigorous convergence result.* Contrary to the 1D case, we will not present the formal analysis allowing to guess that the limit equation is ZK. The principle is indeed identical, but the computations become quite lengthy. Let us refer to [29] for that point.

We state directly the theorem asserting the convergence to ZK:

**Theorem 6.** *Let  $(f_{\varepsilon,0})_{\varepsilon \in (0,1)}$  be a family of non-negative initial data satisfying the assumptions of Theorem 4 and such that there exists  $C > 0$  with:*

$$\mathcal{E}_\varepsilon(0) \leq C, \quad \forall \varepsilon \in (0,1). \quad (3.8)$$

*We denote by  $(f_\varepsilon)_{\varepsilon \in (0,1)}$  a sequence of global weak solutions to (3.1) given by Theorem 4. Let  $\mathcal{H}_\varepsilon$  be the relative entropy defined by the functional:*

$$\begin{aligned} \mathcal{H}_\varepsilon(t) &:= \frac{1}{2} \int f_\varepsilon \left[ |v_1 - u_1^{(1)} - \varepsilon u_1^{(2)}|^2 \right. \\ &\quad \left. + |v_2 - \sqrt{\varepsilon} u_2^{(1)} - \varepsilon u_2^{(2)}|^2 + |v_3 - \sqrt{\varepsilon} u_3^{(1)} - \varepsilon u_3^{(2)}|^2 \right] dv dx \\ &+ \frac{1}{2} \varepsilon \int |\nabla_x \phi_\varepsilon - \nabla_x \phi_1 - \varepsilon \nabla_x \phi_2 - \varepsilon^2 \nabla_x \phi_3|^2 dx \\ &+ \frac{1}{\varepsilon^2} \int \left( e^{\varepsilon \phi_\varepsilon} \log(e^{\varepsilon \phi_\varepsilon} / e^{\varepsilon(\phi_1 + \varepsilon \phi_2 + \varepsilon^2 \phi_3)}) - e^{\varepsilon \phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon \phi_2 + \varepsilon^2 \phi_3)} \right) dx, \end{aligned} \quad (3.9)$$

where

$$\begin{aligned} &(\phi_1, u_1^{(1)}, u_2^{(1)}, u_3^{(1)}, u_1^{(2)}, u_2^{(2)}, u_3^{(2)}, \phi_2, \phi_3) \\ &\in [C([0, T_0[, H^{s+2}(\mathbb{T}^3))]^2 \times [C([0, T_0[, H^{s+1}(\mathbb{T}^3))]^2 \times [C([0, T_0[, H^s(\mathbb{T}^3))]^5 \end{aligned}$$

(with  $s > 3/2 + 1$ ,  $T_0 > 0$ ) satisfy the following system on  $[0, T_0[$ :

$$\begin{cases} 2\partial_t \phi_1 + 2\phi_1 \partial_x \phi_1 + \partial_{x_1}(\Delta + \Delta_\perp)\phi_1 = 0, \\ u_1^{(1)} = \phi_1, \quad u_2^{(1)} = -\partial_{x_3} \phi_1, \quad u_3^{(1)} = \partial_{x_2} \phi_1, \\ \phi_2 = \partial_{x_1 x_1}^2 \phi_1, \\ u_2^{(2)} = \partial_{x_2 x_2}^2 \phi_1, \quad u_3^{(2)} = \partial_{x_3 x_3}^2 \phi_1, \\ \partial_{x_1} u_1^{(2)} = \partial_t u_1 + u_1 \partial_x u_1 + \partial_{x_1 x_1 x_1}^3 \phi_1, \\ \partial_{x_1} \phi_3 = -\partial_t \phi_2 - \partial_{x_1}(u_1^{(2)} \phi_1) - u_1^{(1)} \partial_{x_1} \phi_2 - u_2^{(1)} \partial_{x_2} \phi_2 + u_3^{(1)} \partial_{x_3} \phi_2. \end{cases} \quad (3.10)$$

Then there exist  $C_1, C_2 > 0$ , such that for all  $\varepsilon \in (0, 1)$ ,

$$\forall t \in [0, T_0[, \quad \mathcal{H}_\varepsilon(t) \leq \int_0^t (C_1 \mathcal{H}_\varepsilon(s) ds + C_2 \sqrt{\varepsilon}) ds. \quad (3.11)$$

Assuming in addition that there exists  $C_3$  such that for any  $\varepsilon \in (0, 1)$ :

$$\mathcal{H}_\varepsilon(0) \leq C_3 \sqrt{\varepsilon}, \quad (3.12)$$

then we obtain for any  $\varepsilon \in (0, 1)$ :

$$\forall t \in [0, T_0[, \quad \mathcal{H}_\varepsilon(t) \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}. \quad (3.13)$$

In this theorem, we need a large number of correctors in the relative entropy, precisely because of the nonlinear exponential term in the Poisson equation (compare with the case of Theorem 3, where this term is linearized). In order to get functions satisfying (3.10), we can proceed as follows:

- The existence of  $\phi_1$  is ensured by Theorem 5 (actually the first equation of (3.10) is slightly different from (3.7), but we can come down to (3.7) by using some standard change of variables, see for instance [35]).
- We observe that the six correctors  $(u_2^{(1)}, u_3^{(1)}, u_1^{(2)}, u_2^{(2)}, u_3^{(2)}, \phi_2)$  have their value which is uniquely determined (contrary to the  $1D$  case). We accordingly set:

$$\begin{cases} u_1^{(1)} = \phi_1, \quad u_2^{(1)} = -\partial_{x_3} \phi_1, \quad u_3^{(1)} = \partial_{x_2} \phi_1, \\ \phi_2 = \partial_{x_1 x_1}^2 \phi_1, \\ u_2^{(2)} = \partial_{x_2 x_2}^2 \phi_1, \quad u_3^{(2)} = \partial_{x_3 x_3}^2 \phi_1, \\ \partial_{x_1} u_1^{(2)} = \partial_t u_1 + u_1 \partial_x u_1 + \partial_{x_1 x_1 x_1}^3 \phi_1. \end{cases}$$

- Finally,  $\phi_3$  is a high order corrector whose value is imposed by the last equation of (3.10).

*Remark 4.* Note that a global well-posedness result for ZK in  $\mathbb{T}^3$  would yield global in time estimates (that is  $T_0 = +\infty$ ) in this theorem, which would be significant for logarithmically growing times, as for Theorem 3.

*Remark 5.* From (3.13), we deduce that:

$$\begin{aligned} \frac{1}{\varepsilon^2} \int \left( e^{\varepsilon\phi_\varepsilon} \log(e^{\varepsilon(\phi_\varepsilon)} / e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) - e^{\varepsilon\phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)} \right) dx \\ \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}. \end{aligned} \quad (3.14)$$

Following the terminology in the Boltzmann literature (see for instance the book [41]), this roughly means that  $e^{\varepsilon\phi_\varepsilon}$  “converges entropically” to  $e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}$ .

From the elementary inequality (3.17) (which will be given later), we can also deduce the control:

$$\frac{1}{\varepsilon^2} \int \left( e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - e^{\frac{1}{2}\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)} \right)^2 dx \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}. \quad (3.15)$$

We have as before the corollary:

**Corollary 2.** *Making the same assumptions as in the previous theorem, we obtain the weak convergences:*

$$\begin{cases} \rho_\varepsilon \rightharpoonup_{\varepsilon \rightarrow 0} 1 \text{ in } L_t^\infty \mathcal{M}^1 \text{ weak-}^*, \\ J_\varepsilon \rightharpoonup_{\varepsilon \rightarrow 0} (\phi_1, 0, 0) \text{ in } L_t^\infty \mathcal{M}^1 \text{ weak-}^*. \end{cases} \quad (3.16)$$

Up to some obvious modifications, the proof of Corollary 2 is similar to that of Corollary 1, and therefore we omit it.

*3.3. Proof of Theorem 6.* Relying on the fact the energy  $\mathcal{E}_\varepsilon(t)$  is a non-increasing function of time (and that  $\mathcal{H}_\varepsilon$  is built as a modulation of the energy), we have for all  $t \in [0, T_0[$  and all  $\varepsilon \in (0, 1)$ :

$$\begin{aligned} \mathcal{H}_\varepsilon(t) &\leq \mathcal{H}_\varepsilon(0) + \int_0^t \int \partial_t \left[ f_\varepsilon \left( \frac{1}{2} |u_1^{(1)} + \varepsilon u_1^{(2)}|^2 - v_1 (u_1^{(1)} + \varepsilon u_1^{(2)}) \right) \right] dv dx ds \\ &\quad + \int_0^t \int \partial_t \left[ f_\varepsilon \left( \frac{1}{2} |\sqrt{\varepsilon} u_2^{(1)} + \varepsilon u_2^{(2)}|^2 - v_2 (\sqrt{\varepsilon} u_2^{(1)} + \varepsilon u_2^{(2)}) \right) \right] dv dx ds \\ &\quad + \int_0^t \int \partial_t \left[ f_\varepsilon \left( \frac{1}{2} |\sqrt{\varepsilon} u_3^{(1)} + \varepsilon u_3^{(2)}|^2 - v_3 (\sqrt{\varepsilon} u_3^{(1)} + \varepsilon u_3^{(2)}) \right) \right] dv dx ds \\ &\quad + \varepsilon \int_0^t \int \partial_t \left[ \frac{1}{2} |\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2|^2 - \nabla_x \phi_\varepsilon \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2) \right] dx ds \\ &\quad + \frac{1}{\varepsilon^2} \int_0^t \int \partial_t \left[ e^{\varepsilon\phi_\varepsilon} \log(1/e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)} \right] dx ds \\ &:= \mathcal{H}_\varepsilon(0) + \int_0^t (I_1^1 + I_1^2 + I_1^3 + I_2 + I_3) ds. \end{aligned}$$

The general strategy in the proof will be to keep (without making approximation) all dangerous modulated terms of the form

$$\varepsilon^\alpha \int f_\varepsilon \left( \begin{array}{l} u_1^{(1)} + \varepsilon u_1^{(2)} - v_1 \\ \sqrt{\varepsilon} u_2^{(1)} + \varepsilon u_2^{(2)} - v_2 \\ \sqrt{\varepsilon} u_3^{(1)} + \varepsilon u_3^{(2)} - v_3 \end{array} \right) [\dots] dv dx$$

and

$$\varepsilon^\alpha \int (-e^{\varepsilon\phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) [\dots] dx,$$

where [...] contains some expression independent of  $\varepsilon$ , as soon as  $\alpha \leq 0$ . Then (3.10) is precisely designed so that all terms exactly vanish in the end.

On the contrary, for  $\alpha > 0$ , these terms will be of order  $\varepsilon^\alpha$  (and thus small). For the first type of terms, this can be seen with the same argument as in the proof of theorem 3, namely the uniform (in  $\varepsilon$ ) bounds on the  $L^1$  norm of  $\rho_\varepsilon$  and  $J_\varepsilon$ . For the second type of terms, one has to use the Poisson equation satisfied by  $\phi_\varepsilon$  and use the bound on the electric energy. Indeed, given some smooth function  $\Psi$ , we can write:

$$\varepsilon^\alpha \int e^{\varepsilon\phi_\varepsilon} \Psi dx = \varepsilon^\alpha \int \varepsilon^2 \Delta \phi_\varepsilon \Psi dx + \varepsilon^\alpha \int \rho_\varepsilon \Psi dx.$$

The second term is clearly of order  $\varepsilon^\alpha$ , using the uniform  $L^1$  bound on  $\rho_\varepsilon$ . On the other hand, by integration by parts, we have for the first term

$$\varepsilon^\alpha \int \varepsilon^2 \Delta \phi_\varepsilon \Psi dx = \varepsilon^{\alpha+2} \int \nabla \phi_\varepsilon \cdot \nabla \Psi dx,$$

which is of order  $\varepsilon^{\alpha+1}$  using the uniform bound obtained thanks to the conservation of energy:

$$\varepsilon \int |\nabla \phi_\varepsilon|^2 dx \leq \mathcal{E}_\varepsilon(0) \leq C.$$

Finally any term like  $\varepsilon^\alpha \int e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)} [\dots] dx$  is clearly of order  $\varepsilon^\alpha$ .

We start by studying separately  $I_1^1 + I_1^2 + I_1^3$ ,  $I_2$  and  $I_3$ . All computations are justified using only the local conservation laws (3.5) and (3.6).

**Study of  $I_1^1 + I_1^2 + I_1^3$ .**

With similar computations as in the proof of Theorem 3, we obtain the identity

$$\begin{aligned}
I_1^1 + I_1^2 + I_1^3 = & \\
& \frac{1}{\varepsilon} \int f_\varepsilon \begin{pmatrix} u_1^{(1)} + \varepsilon u_1^{(2)} - v_1 \\ \sqrt{\varepsilon} u_2^{(1)} + \varepsilon u_2^{(2)} - v_2 \\ \sqrt{\varepsilon} u_3^{(1)} + \varepsilon u_3^{(2)} - v_3 \end{pmatrix} \cdot \begin{pmatrix} -\partial_{x_1} u_1^{(1)} + \partial_{x_1} \phi_1 \\ -u_3^{(1)} + \partial_{x_2} \phi_1 \\ u_2^{(1)} + \partial_{x_3} \phi_1 \end{pmatrix} dv dx \\
& - \int \frac{1}{\varepsilon} \rho_\varepsilon u^{(1)} \cdot \nabla_x \phi_1 dx - \int \rho_\varepsilon u^{(2)} \cdot \nabla_x \phi_1 dx + \int \frac{1}{\varepsilon} J_\varepsilon \cdot \nabla_x \phi_1 dx \\
& + \frac{1}{\sqrt{\varepsilon}} \int f_\varepsilon \begin{pmatrix} \sqrt{\varepsilon} u_2^{(1)} + \varepsilon u_2^{(2)} - v_2 \\ \sqrt{\varepsilon} u_3^{(1)} + \varepsilon u_3^{(2)} - v_3 \end{pmatrix} \cdot \begin{pmatrix} -u_3^{(2)} - \partial_{x_1} u_2^{(1)} \\ u_2^{(2)} - \partial_{x_1} u_3^{(1)} \end{pmatrix} dv dx \\
& + \int f_\varepsilon \begin{pmatrix} u_1^{(1)} + \varepsilon u_1^{(2)} - v_1 \\ \sqrt{\varepsilon} u_2^{(1)} + \varepsilon u_2^{(2)} - v_2 \\ \sqrt{\varepsilon} u_3^{(1)} + \varepsilon u_3^{(2)} - v_3 \end{pmatrix} \cdot \begin{pmatrix} \partial_t u_1^{(1)} + u_1^{(1)} \partial_{x_1} u_1^{(1)} - \partial_{x_1} u_2^{(1)} + \partial_{x_1} \phi_2 \\ -\partial_{x_1} u_2^{(2)} + \partial_{x_2} \phi_2 \\ -\partial_{x_1} u_3^{(2)} + \partial_{x_3} \phi_2 \end{pmatrix} dv dx \\
& + \int J_\varepsilon \cdot \nabla_x \phi_2 dx - \int \rho_\varepsilon u^{(1)} \cdot \nabla_x \phi_2 dx - \varepsilon \int \rho_\varepsilon u^{(2)} \cdot \nabla_x \phi_2 dx \\
& - \int f_\varepsilon (u_1^{(1)} + \varepsilon u_1^{(2)} - v_1)^2 \partial_x u_1^{(1)} dx \\
& - \int \frac{1}{\varepsilon} \rho_\varepsilon E_\varepsilon \cdot u^{(1)} dx - \int \rho_\varepsilon E_\varepsilon \cdot u^{(2)} dx \\
& + \mathcal{O}(\varepsilon).
\end{aligned}$$

In this equation,  $\mathcal{O}(\varepsilon)$  is as usual a notation for all terms that can be bounded by  $C\varepsilon$ , where  $C$  is a positive constant. We also denote here:

$$u^{(1)} := (u_1^{(1)}, \sqrt{\varepsilon} u_2^{(1)}, \sqrt{\varepsilon} u_3^{(1)}) \text{ and } u^{(2)} := (u_1^{(2)}, u_2^{(2)}, u_3^{(2)}).$$

We observe that we can write:

$$\begin{aligned}
& - \int \frac{1}{\varepsilon} \rho_\varepsilon E_\varepsilon \cdot u^{(1)} dx - \int \frac{1}{\varepsilon} \rho_\varepsilon u^{(1)} \cdot \nabla_x \phi_1 dx - \int \rho_\varepsilon u^{(1)} \cdot \nabla_x \phi_2 dx \\
& \quad = -\frac{1}{\varepsilon} \int \rho_\varepsilon u_1 \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 - \nabla_x \phi_\varepsilon).
\end{aligned}$$

### Study of $I_2$ .

We obtain:

$$\begin{aligned}
I_2 = & \varepsilon \int \partial_t (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3) \cdot (-\nabla_x \phi_\varepsilon + \nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3) dx \\
& - \varepsilon \int \partial_t \nabla_x \phi_\varepsilon \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3) dx.
\end{aligned}$$

We have the easy bound:

$$\begin{aligned} & \varepsilon \left| \int \partial_t (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3) \cdot (-\nabla_x \phi_\varepsilon + \nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3) dx \right| \\ & \leq C \left[ \varepsilon \int (\partial_t (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3))^2 dx + \varepsilon \int (-\partial_x \phi_\varepsilon + \partial_x \phi_1 + \varepsilon \partial_x \phi_2)^2 dx \right] \\ & \leq \mathcal{O}(\varepsilon) + C\mathcal{H}_\varepsilon(t). \end{aligned}$$

### Study of $I_3$ .

We can compute:

$$\begin{aligned} I_3 &= \frac{1}{\varepsilon} \int (-e^{\varepsilon\phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) \partial_t (\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3) dx \\ & \quad - \frac{1}{\varepsilon} \int \partial_t (e^{\varepsilon\phi_\varepsilon}) (\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3) dx \\ &= I_3^1 + I_3^2. \end{aligned}$$

The first term can be recast as follows:

$$\begin{aligned} I_3^1 &= \frac{1}{\varepsilon} \int (-e^{\varepsilon\phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) \partial_t \phi_1 dx + \int (-e^{\varepsilon\phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) \partial_t \phi_2 dx \\ & \quad + \varepsilon \int (-e^{\varepsilon\phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) \partial_t \phi_3 dx. \end{aligned}$$

For the second one, we use the Poisson equation (with a time derivative) and the conservation of charge in shifted variables, that we recall below:

$$\begin{cases} \partial_t (e^{\varepsilon\phi_\varepsilon}) = \varepsilon^2 \Delta \partial_t \phi_\varepsilon + \partial_t \rho_\varepsilon, \\ \frac{1}{\varepsilon} \partial_t \rho_\varepsilon = \frac{1}{\varepsilon^2} \partial_{x_1} \rho_\varepsilon - \frac{1}{\varepsilon} \nabla_x \cdot J_\varepsilon, \end{cases}$$

to obtain:

$$\begin{aligned} I_3^2 &= -\frac{1}{\varepsilon^2} \int \partial_{x_1} \rho_\varepsilon (\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3) dx \\ & \quad + \frac{1}{\varepsilon} \int \nabla_x \cdot J_\varepsilon \phi_1 dx + \int \nabla_x \cdot J_\varepsilon \phi_2 dx + \varepsilon \int \nabla_x \cdot J_\varepsilon \phi_3 dx \\ & \quad - \varepsilon \int \partial_t \Delta \phi_\varepsilon (\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3) dx. \end{aligned}$$

As in the proof of Theorem 3, every term in  $I_3^2$  but the first one gets simplified with some other ones obtained in  $I_1$  and  $I_2$ .

### Study of the remaining significant terms.



There remain to study the following potentially harmful terms:

$$\begin{aligned}
K_1 &:= - \int \rho_\varepsilon u^{(2)} \cdot \nabla_x (\phi_1 + \varepsilon \phi_2) dx, & K_2 &:= - \int \rho_\varepsilon E_\varepsilon \cdot u^{(2)} dx, \\
K_3 &:= - \frac{1}{\varepsilon} \int \rho_\varepsilon u^{(1)} \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 - \nabla_x \phi_\varepsilon) dx, \\
K_4 &:= - \frac{1}{\varepsilon^2} \int \partial_{x_1} \rho_\varepsilon (\phi_1 + \varepsilon \phi_2 + \varepsilon^2 \phi_3) dx, \\
K_5 &:= \frac{1}{\varepsilon} \int (-e^{\varepsilon \phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon \phi_2 + \varepsilon^2 \phi_3)}) \partial_t \phi_1 dx. \\
K_6 &:= \int (-e^{\varepsilon \phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon \phi_2 + \varepsilon^2 \phi_3)}) \partial_t \phi_2 dx.
\end{aligned}$$

We start with  $K_1 + K_2$ , using the Poisson equation:

$$\begin{aligned}
K_1 + K_2 &= - \int e^{\varepsilon \phi_\varepsilon} u^{(2)} \cdot \nabla_x (\phi_1 + \varepsilon \phi_2 - \phi_\varepsilon) dx \\
&\quad + \varepsilon^2 \int \Delta \phi_\varepsilon u^{(2)} \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 - \nabla_x \phi_\varepsilon) dx. \\
&= L_1 + L_2.
\end{aligned}$$

We have:

$$\begin{aligned}
L_1 &= - \int e^{\varepsilon \phi_\varepsilon} u^{(2)} \cdot \nabla_x \phi_1 dx - \varepsilon \int e^{\varepsilon \phi_\varepsilon} u^{(2)} \cdot \nabla_x \phi_2 dx \\
&\quad - \frac{1}{\varepsilon} \int e^{\varepsilon \phi_\varepsilon} \nabla_x \cdot u^{(2)} dx.
\end{aligned}$$

For  $L_2$ , we have

$$\begin{aligned}
L_2 &= \varepsilon^2 \int \Delta \phi_\varepsilon u^{(2)} \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2) dx - \varepsilon^2 \int \Delta \phi_\varepsilon u^{(2)} \cdot \nabla_x \phi_\varepsilon dx \\
&:= L_2^1 + L_2^2.
\end{aligned}$$

By integration by parts, we get:

$$\begin{aligned}
|L_2^1| &= \left| \varepsilon^2 \int \nabla \phi_\varepsilon \cdot u^{(2)} (\Delta \phi_1 + \varepsilon \Delta \phi_2) dx \right| \\
&\leq C \left( \varepsilon^2 \int |\nabla \phi_\varepsilon|^2 dx + C\varepsilon \right) \\
&\leq C(\varepsilon \mathcal{E}_\varepsilon(t) + C\varepsilon) \leq C\varepsilon.
\end{aligned}$$

Note here that we have used the Lipschitz bound on the second derivative of  $\phi_1$  and  $\phi_2$ .

For  $L_2^2$ , we rely on the usual trick to write:

$$\begin{aligned}
L_2^2 &= - \frac{1}{2} \varepsilon^2 \int \nabla_x |\nabla_x \phi_\varepsilon|^2 \cdot u^{(2)} dx \\
&= \frac{1}{2} \varepsilon^2 \int |\nabla_x \phi_\varepsilon|^2 \nabla_x \cdot u^{(2)} dx,
\end{aligned}$$

from which we deduce that

$$|L_2^2| \leq C\varepsilon.$$

Using the Poisson equation for  $K_3$ :

$$\begin{aligned} K_3 &= -\frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} u^{(1)} \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 - \nabla_x \phi_\varepsilon) dx \\ &\quad + \int \varepsilon \Delta \phi_\varepsilon u^{(1)} \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 - \nabla_x \phi_\varepsilon) dx \\ &:= K_3^1 + K_3^2. \end{aligned}$$

Concerning the first term, we write the decomposition:

$$\begin{aligned} K_3^1 &= -\frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} u^{(1)} \cdot \nabla_x \phi_1 dx \\ &\quad - \int e^{\varepsilon\phi_\varepsilon} u^{(1)} \cdot \nabla_x \phi_2 dx + \frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} u^{(1)} \cdot \nabla_x \phi_\varepsilon dx \\ &:= J_1 + J_2 + J_3. \end{aligned}$$

We start by the study of  $J_1$ :

$$\begin{aligned} J_1 &= \frac{1}{\varepsilon} \int \left( e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)} - e^{\varepsilon\phi_\varepsilon} \right) u_1^{(1)} \partial_{x_1} \phi_1 dx \\ &\quad - \frac{1}{\varepsilon} \int e^{\varepsilon(\phi_1 - \varepsilon\phi_2)} u_1^{(1)} \partial_{x_1} \phi_1 dx - \frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} (u^{(1)} - u_1^{(1)} e_1) \cdot \nabla_x \phi_1 dx. \end{aligned}$$

Clearly, we have (for instance using a Taylor inequality):

$$-\frac{1}{\varepsilon} \int e^{\varepsilon(\phi_1 - \varepsilon\phi_2)} u_1^{(1)} \partial_{x_1} \phi_1 dx = -\frac{1}{\varepsilon} \int u_1^{(1)} \partial_{x_1} \phi_1 dx + \int \phi_1 u_1^{(1)} \partial_{x_1} \phi_1 dx + \mathcal{O}(\varepsilon).$$

In the end, we will take  $u_1^{(1)} = \phi_1$ , so the terms of this latest expression which are of order  $\mathcal{O}(1/\varepsilon)$  and  $\mathcal{O}(1)$  are exactly equal to 0.

For  $J_2$ , it is sufficient to write:

$$J_2 = - \int e^{\varepsilon\phi_\varepsilon} u_1^{(1)} \partial_{x_1} \phi_2 dx - \int e^{\varepsilon\phi_\varepsilon} (u^{(1)} - u_1^{(1)} e_1) \cdot \nabla_x \phi_2 dx.$$

We decompose the last term in the following way

$$\begin{aligned} J_3 &= \frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} \phi_1 \partial_{x_1} \phi_\varepsilon \\ &\quad + \frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} (u_1^{(1)} - \phi_1) \partial_{x_1} \phi_\varepsilon \\ &\quad + \frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} (u^{(1)} - u_1^{(1)} e_1) \cdot \nabla_x \phi_\varepsilon. \end{aligned}$$

The first term could be dangerous, but will disappear using a term coming from a term of  $K_4$  (first term of  $K_4^1$  below). The second one will be equal to 0 since we take  $u_1^{(1)} = \phi_1$ .

Gathering the pieces together, we finally obtain:

$$\begin{aligned} \frac{1}{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} (u^{(1)} - u_1^{(1)} e_1) \cdot \nabla_x (\phi_\varepsilon - \phi_1 - \varepsilon\phi_2) = \\ - \frac{1}{\varepsilon^{3/2}} \int e^{\varepsilon\phi_\varepsilon} (\partial_{x_2} u_2^{(1)} + \partial_{x_3} u_3^{(1)}) dx \\ - \frac{1}{\sqrt{\varepsilon}} \int e^{\varepsilon\phi_\varepsilon} (u_2^{(1)} \partial_{x_2} \phi_1 + u_3^{(1)} \partial_{x_3} \phi_1) dx \\ - \sqrt{\varepsilon} \int e^{\varepsilon\phi_\varepsilon} (u_2^{(1)} \partial_{x_2} \phi_2 + u_3^{(1)} \partial_{x_3} \phi_2) dx. \end{aligned}$$

Let us now turn to the treatment of  $K_3^2$ :

$$K_3^2 = \int \varepsilon \Delta \phi_\varepsilon u^{(1)} \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2) dx - \varepsilon/2 \int \nabla_x |\nabla_x \phi_\varepsilon|^2 \cdot u^{(1)} dx.$$

This term is treated exactly as  $L_2$ , but we have to be careful since it is singular in  $\varepsilon$ . Here, rather than bounding by the energy, we shall rely on a bound by the modulated energy. We have

$$\begin{aligned} \varepsilon/2 \int \nabla_x |\nabla_x \phi_\varepsilon|^2 \cdot u^{(1)} dx \\ = -\varepsilon/2 \int |\nabla_x \phi_\varepsilon|^2 \nabla_x \cdot u^{(1)} dx \\ - \varepsilon/2 \int |\nabla_x \phi_\varepsilon - \nabla_x \phi_1 - \varepsilon \nabla_x \phi_2 - \varepsilon^2 \nabla_x \phi_3|^2 \nabla_x \cdot u^{(1)} dx \\ + \varepsilon/2 \int |\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3|^2 \nabla_x \cdot u^{(1)} dx \\ - \varepsilon \int \nabla_x \phi_\varepsilon \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3) \nabla_x \cdot u^{(1)} dx. \end{aligned}$$

We shall bound the very last term using the Cauchy-Schwarz inequality and the various Lipschitz bounds:

$$\begin{aligned} \left| \varepsilon \int \nabla_x \phi_\varepsilon \cdot (\nabla_x \phi_1 + \varepsilon \nabla_x \phi_2 + \varepsilon^2 \nabla_x \phi_3) \nabla_x \cdot u^{(1)} dx \right| \\ \leq C\sqrt{\varepsilon} \sqrt{\varepsilon \int |\nabla_x \phi_\varepsilon|^2 dx} \leq C\sqrt{\varepsilon} \mathcal{E}_\varepsilon(t) \leq C\sqrt{\varepsilon}. \end{aligned}$$

Therefore we obtain:

$$|K_3^2| \leq C\mathcal{H}_\varepsilon(t) + C\sqrt{\varepsilon}.$$

Finally, concerning  $K_4$  we get:

$$\begin{aligned} K_4 &= \frac{1}{\varepsilon^2} \int \rho_\varepsilon (\partial_{x_1} \phi_1 + \varepsilon \partial_{x_1} \phi_2 + \varepsilon^2 \partial_{x_1} \phi_3) dx \\ &= \frac{1}{\varepsilon^2} \int e^{\varepsilon \phi_\varepsilon} (\partial_{x_1} \phi_1 + \varepsilon \partial_{x_1} \phi_2 + \varepsilon^2 \partial_{x_1} \phi_3) dx \\ &\quad - \int \Delta \phi_\varepsilon (\partial_{x_1} \phi_1 + \varepsilon \partial_{x_1} \phi_2 + \varepsilon^2 \partial_{x_1} \phi_3) dx \\ &:= K_4^1 + K_4^2. \end{aligned}$$

The first term can be decomposed as:

$$\begin{aligned} K_4^1 &= -\frac{1}{\varepsilon} \int \partial_{x_1} \phi_\varepsilon e^{\varepsilon \phi_\varepsilon} \phi_1 dx \\ &\quad + \frac{1}{\varepsilon} \int e^{\varepsilon \phi_\varepsilon} \partial_{x_1} \phi_2 dx \\ &\quad + \int e^{\varepsilon \phi_\varepsilon} \partial_{x_1} \phi_3 dx. \end{aligned}$$

The second one can be recast as:

$$K_4^2 = - \int \phi_\varepsilon \Delta \partial_{x_1} \phi_1 dx + \mathcal{O}(\sqrt{\varepsilon}).$$

We shall focus our attention on the important term:

$$\begin{aligned} L &:= - \int \phi_\varepsilon \Delta \partial_{x_1} \phi_1 dx \\ &= -\frac{1}{\varepsilon} \int e^{\varepsilon \phi_\varepsilon} \Delta \partial_{x_1} \phi_1 dx + \frac{1}{\varepsilon} \int (e^{\varepsilon \phi_\varepsilon} - \varepsilon \phi_\varepsilon) \Delta \partial_{x_1} \phi_1 dx. \end{aligned}$$

We have the following technical result, allowing to consider the second term above as an error term:

**Lemma 1.** *There exists  $C > 0$  such that for any  $\varepsilon \in (0, 1)$ :*

$$\frac{1}{\varepsilon} \left| \int (e^{\varepsilon \phi_\varepsilon} - \varepsilon \phi_\varepsilon) \Delta \partial_{x_1} \phi_1 dx \right| \leq C \sqrt{\varepsilon}.$$

*Proof (Proof of Lemma 1).*

The naive idea would be to use the Taylor expansion

$$e^x \sim_0 1 + x + \frac{1}{2} x^2.$$

but we cannot say that  $\varepsilon \|\phi_\varepsilon\|_\infty \ll 1$ . (Even worse, we do not have any  $L^2$  control on  $\phi_\varepsilon$ .) Instead, we will only rely on the bounds given by the conservation of energy. The classical inequality (valid for  $x, y > 0$ ) will be very useful:

$$(\sqrt{x} - \sqrt{y})^2 \leq x \log(x/y) - x + y. \quad (3.17)$$

We shall write the decomposition

$$\begin{aligned} \frac{1}{\varepsilon} \int (e^{\varepsilon\phi_\varepsilon} - \varepsilon\phi_\varepsilon) \Delta \partial_{x_1} \phi_1 dx &= \\ &= \frac{1}{\varepsilon} \int (e^{\varepsilon\phi_\varepsilon} - 2e^{\frac{1}{2}\varepsilon\phi_\varepsilon}) \Delta \partial_{x_1} \phi_1 dx \\ &\quad + \frac{1}{\varepsilon} \int (e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - \frac{1}{2}\varepsilon\phi_\varepsilon) 2\Delta \partial_{x_1} \phi_1 dx. \end{aligned}$$

We first recast the first term as follows:

$$\begin{aligned} \frac{1}{\varepsilon} \int (e^{\varepsilon\phi_\varepsilon} - 2e^{\frac{1}{2}\varepsilon\phi_\varepsilon}) \Delta \partial_{x_1} \phi_1 dx &= \frac{1}{\varepsilon} \int (e^{\varepsilon\phi_\varepsilon} - 2e^{\frac{1}{2}\varepsilon\phi_\varepsilon} + 1) \Delta \partial_{x_1} \phi_1 dx \\ &= \frac{1}{\varepsilon} \int (e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - 1)^2 \Delta \partial_{x_1} \phi_1 dx. \end{aligned}$$

Using (3.17) and the Lipschitz bound on the second order derivative of  $\phi_1$ , we obtain:

$$\begin{aligned} \frac{1}{\varepsilon} \int (e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - 1)^2 \Delta \partial_{x_1} \phi_1 dx &\leq C\varepsilon \int \frac{1}{\varepsilon^2} (e^{\varepsilon\phi_\varepsilon} (\varepsilon\phi_\varepsilon - 1) - 1) dx \\ &\leq \varepsilon \mathcal{E}_\varepsilon(t) \\ &\leq \varepsilon \mathcal{E}_\varepsilon(0) \\ &\leq C\varepsilon. \end{aligned}$$

For the second term, we have by integration by parts:

$$\frac{1}{\varepsilon} \int (e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - \frac{1}{2}\varepsilon\phi_\varepsilon) 2\Delta \partial_{x_1} \phi_1 dx = - \int \nabla_x \phi_\varepsilon (e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - 1) \cdot \nabla_x \partial_{x_1} \phi_1 dx.$$

Then we write, using  $|ab| \leq \frac{1}{2}(\lambda a^2 + \frac{1}{\lambda} b^2)$ , with  $\alpha$  a parameter to be fixed:

$$\begin{aligned} \left| \int \nabla_x \phi_\varepsilon (e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - 1) \cdot \nabla_x \partial_{x_1} \phi_1 dx \right| \\ \leq C\varepsilon^{\alpha-1} \int |\nabla_x \phi_\varepsilon|^2 + C\varepsilon^{-\alpha+2} \int \frac{1}{\varepsilon^2} (e^{\varepsilon\phi_\varepsilon} (\varepsilon\phi_\varepsilon - 1) + 1) dx. \end{aligned}$$

To optimize, it is clear that we should take  $\alpha = 3/2$ . Thus, relying on the uniform bound given by the energy, we get:

$$\left| \frac{1}{\varepsilon} \int \left( e^{\frac{1}{2}\varepsilon\phi_\varepsilon} - \frac{1}{2}\varepsilon\phi_\varepsilon \right) 2\Delta \partial_{x_1} \phi_1 dx \right| \leq C\sqrt{\varepsilon}.$$

The proof is therefore complete.

**Conclusion.**

Finally, we impose all cancellations for the terms

$$\varepsilon^\alpha \int f_\varepsilon \left( \begin{array}{l} u_1^{(1)} + \varepsilon u_1^{(2)} - v_1 \\ \sqrt{\varepsilon} u_2^{(1)} + \varepsilon u_2^{(2)} - v_2 \\ \sqrt{\varepsilon} u_3^{(1)} + \varepsilon u_3^{(2)} - v_3 \end{array} \right) [\dots] dv dx$$

for  $\alpha = -1, -1/2, 0$  and

$$\varepsilon^\beta \int (-e^{\varepsilon\phi_\varepsilon} + e^{\varepsilon(\phi_1 + \varepsilon\phi_2 + \varepsilon^2\phi_3)}) [\dots] dx$$

for  $\beta = -3/2, -1, -1/2, 0$ , which precisely means that we impose:

$$\left\{ \begin{array}{l} -\partial_{x_1} u_1^{(1)} + \partial_{x_1} \phi_1 = 0, \\ -u_3^{(1)} + \partial_{x_2} \phi_1 = 0, \\ u_2^{(1)} + \partial_{x_3} \phi_1 = 0, \\ -u_3^{(2)} - \partial_{x_1} u_2^{(1)} = 0, \\ u_2^{(2)} - \partial_{x_1} u_3^{(1)} = 0, \\ \partial_t u_1^{(1)} + u_1^{(1)} \partial_{x_1} u_1^{(1)} - \partial_{x_1} u_2^{(1)} + \partial_{x_1} \phi_2 = 0, \\ -\partial_{x_1} u_2^{(2)} + \partial_{x_2} \phi_2 = 0, \\ -\partial_{x_1} u_3^{(2)} + \partial_{x_3} \phi_2 = 0, \\ \partial_{x_2} u_2^{(1)} + \partial_{x_3} u_3^{(1)} = 0, \\ \partial_t \phi_1 + \nabla_x \cdot u^{(2)} + \phi_1 \partial_{x_1} \phi_1 + \Delta \partial_{x_1} \phi_1 - \partial_{x_1} \phi_2 = 0, \\ u_2^{(1)} \partial_{x_2} \phi_1 + u_3^{(1)} \partial_{x_3} \phi_1 = 0, \\ \partial_t \phi_2 + \phi_1 \partial_{x_1} \phi_2 + u_1^{(2)} \partial_{x_1} \phi_1 + u_2^{(2)} \partial_{x_2} \phi_1 + u_3^{(2)} \partial_{x_3} \phi_1 - \partial_{x_1} \phi_3 = 0. \end{array} \right. \quad (3.18)$$

All these identities are exactly obtained as a consequence of (3.10).

We end up with:

$$\mathcal{H}_\varepsilon(t) \leq \mathcal{H}_\varepsilon(0) + \int_0^t (C_1 \mathcal{H}_\varepsilon(s) + C_2 \sqrt{\varepsilon}) ds, \quad (3.19)$$

and arguing as in the proof of Theorem 3, this is over.

#### 4. Appendix A: From the Vlasov-Poisson equation to the Kadomstev-Petviashvili II equation

Assuming a slow variation in the  $x_2$  direction, one may end up with the following anisotropic long wave scaling for the Vlasov-Poisson system for ions. For simplicity, we restrict here to the linearized Maxwell-Boltzmann law, but the same study

can be performed for the full equations. The variables are  $t \geq 0, x \in \mathbb{T}^2, v \in \mathbb{R}^2$ :

$$\begin{cases} \varepsilon \partial_t f_\varepsilon - \partial_{x_1} f_\varepsilon + \varepsilon v_1 \partial_{x_1} f_\varepsilon + \varepsilon^{3/2} v_2 \partial_{x_2} f_\varepsilon + E_\varepsilon \cdot \nabla_v f_\varepsilon = 0, \\ E = (-\partial_{x_1} \phi_\varepsilon, -\sqrt{\varepsilon} \partial_{x_2} \phi_\varepsilon), \\ -\varepsilon^2 \partial_{x_1 x_1}^2 \phi_\varepsilon - \varepsilon^3 \partial_{x_2 x_2}^2 \phi_\varepsilon + \varepsilon \phi_\varepsilon = \int_{\mathbb{R}^2} f_\varepsilon dv - 1, \\ f|_{t=0} = f_0. \end{cases} \quad (4.1)$$

The scaled energy of this system is the following functional:

$$\begin{aligned} \mathcal{E}_\varepsilon(t) &:= \frac{1}{2} \int f_\varepsilon |v|^2 dv dx \\ &+ \frac{1}{2} \varepsilon \int |\partial_{x_1} \phi_\varepsilon|^2 dx + \frac{1}{2} \varepsilon^2 \int |\partial_{x_2} \phi_\varepsilon|^2 dx + \frac{1}{2} \int \phi_\varepsilon^2 dx. \end{aligned} \quad (4.2)$$

We have the existence of global weak solutions, sharing the same properties with those given in Theorem 1.

For the KP-II equation, that is

$$\partial_{x_1} (\partial_t \phi_1 + \phi_1 \partial_{x_1} \phi_1 + \partial_{x_1 x_1 x_1}^3 \phi_1) + \partial_{x_2 x_2}^2 \phi_1 = 0, \quad (4.3)$$

our reference is an article by Bourgain [12], in which is proved the following theorem:

**Theorem 7.** *The KP-II equation is globally well-posed in  $H^s(\mathbb{T}^2)$  for  $s \geq 0$ .*

Once again, we will only use this theorem for large values of  $s$ .

As for the other cases, we obtain the rigorous convergence to KP-II, which is summarized in the following theorem:

**Theorem 8.** *Let  $(f_{\varepsilon,0})_{\varepsilon \in (0,1)}$  be a family of non-negative initial data such that*

$$\|f_{\varepsilon,0}\|_{L^1 \cap L^\infty} < +\infty, \quad \int f_{\varepsilon,0} dv dx = 1, \quad (4.4)$$

and such that there exists  $C > 0$  with:

$$\mathcal{E}_\varepsilon(0) \leq C, \quad \forall \varepsilon \in (0,1). \quad (4.5)$$

We denote by  $(f_\varepsilon)_{\varepsilon \in (0,1)}$  a sequence of global weak solutions to (4.1). Let  $\mathcal{H}_\varepsilon$  be the relative entropy defined by the functional:

$$\begin{aligned} \mathcal{H}_\varepsilon(t) &:= \frac{1}{2} \int f_\varepsilon \left[ |v_1 - u_1^{(1)} - \varepsilon u_1^{(2)}|^2 + |v_2 - \sqrt{\varepsilon} u_2^{(1)} - \varepsilon^{3/2} u_2^{(2)}|^2 \right] dv dx \\ &+ \frac{1}{2} \varepsilon \int |\partial_{x_1} \phi_\varepsilon - \partial_{x_1} \phi_1 - \varepsilon \partial_{x_1} \phi_2|^2 dx + \frac{1}{2} \varepsilon^2 \int |\partial_{x_2} \phi_\varepsilon - \partial_{x_2} \phi_1 - \varepsilon \partial_{x_2} \phi_2|^2 dx \\ &+ \frac{1}{2} \int (\phi_\varepsilon - \phi_1 - \varepsilon \phi_2)^2 dx, \end{aligned} \quad (4.6)$$

where  $(u_1, u_2, \phi_1, \phi_2) \in [C([0, +\infty[, H^{s+2}(\mathbb{T}^2))]^4$  (with  $s > 2$ ) satisfy the following system:

$$\begin{cases} \partial_{x_1} (2\partial_t \phi_1 + 3\phi_1 \partial_{x_1} \phi_1 + \partial_{x_1 x_1 x_1}^3 \phi_1) + \partial_{x_2 x_2}^2 \phi_1 = 0, \\ u_1^{(1)} = \phi_1, \\ \partial_{x_1} u_2^{(1)} = \partial_{x_2} \phi_1, \\ \partial_{x_1} (u_1^{(2)} - \phi_2) = \partial_t \phi_1 + \phi_1 \partial_{x_1} \phi_1. \end{cases} \quad (4.7)$$

Then there exist  $C_1, C_2 > 0$ , such that for all  $\varepsilon \in (0, 1)$ ,

$$\mathcal{H}_\varepsilon(t) \leq \mathcal{H}_\varepsilon(0) + \int_0^t (C_1 \mathcal{H}_\varepsilon(s) ds + C_2 \sqrt{\varepsilon}) ds. \quad (4.8)$$

Assuming in addition that there exists  $C_3 > 0$  such that for any  $\varepsilon \in (0, 1)$ ,

$$\mathcal{H}_\varepsilon(0) \leq C_3 \sqrt{\varepsilon}. \quad (4.9)$$

Then we obtain for all  $\varepsilon \in (0, 1)$ :

$$\forall t \in [0, +\infty[, \quad \mathcal{H}_\varepsilon(t) \leq C_3 e^{C_1 t} \sqrt{\varepsilon} + C_2 \frac{e^{C_1 t} - 1}{C_1} \sqrt{\varepsilon}, \quad (4.10)$$

as well as the weak convergences:

$$\begin{cases} \rho_\varepsilon \rightharpoonup_{\varepsilon \rightarrow 0} 1 \text{ in } L_t^\infty \mathcal{M}^1 \text{ weak-}^*, \\ J_\varepsilon \rightharpoonup_{\varepsilon \rightarrow 0} (\phi_1, 0) \text{ in } L_t^\infty \mathcal{M}^1 \text{ weak-}^*. \end{cases} \quad (4.11)$$

The proof of Theorem 8 follows from computations in the same spirit as the previous ones, and therefore we leave it to the reader.

## 5. Appendix B: A KdV limit in the whole space $\mathbb{R}$

All results stated in this paper are restricted to PDEs set in the torus for the space variable. The reason is that in all cases, in the end, the first moment  $\rho_\varepsilon$  (charge density) has to weakly converge to the constant 1. This function is obviously not integrable in the whole space, and the assumptions needed for our results to hold are actually not consistent in the whole space case.

It is nevertheless possible to slightly adapt the KdV limit of Section 2 to handle that case. Keeping the same notations, we shall rely on the fact that the electric potential can be defined up to a constant, and use the following version of the energy

$$\mathcal{F}_\varepsilon(t) := \frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} f_\varepsilon |v|^2 dv dx + \frac{1}{2} \varepsilon \int_{\mathbb{R}} |\nabla_x \phi_\varepsilon|^2 dx + \frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} \left( \phi_\varepsilon - \frac{1}{\varepsilon} \right)^2 dx dx, \quad (5.1)$$

This means that  $\phi_\varepsilon \in \frac{1}{\varepsilon} + L^2(\mathbb{R})$  (instead of the more usual  $\phi_\varepsilon \in L^2(\mathbb{R})$ ).



The Ansatz in that case for the formal computations now corresponds to the following one:

$$\begin{cases} \rho_\varepsilon = \varepsilon\rho_1 + \mathcal{O}(\varepsilon^2), \\ \phi_\varepsilon = \frac{1}{\varepsilon} + \phi_1 + \varepsilon\phi_2 + \mathcal{O}(\varepsilon^2), \\ u_\varepsilon = u_1 + \varepsilon u_2 + \mathcal{O}(\varepsilon^2). \end{cases} \quad (5.2)$$

In particular this means that we expect that  $\rho_\varepsilon$  weakly converges to 0 (which is of course integrable on  $\mathbb{R}$ ). The formal computations then remain the same. In the end, we may obtain the same result as Theorem 3 except that we consider the following relative entropy instead of (2.21):

$$\begin{aligned} \mathcal{H}_\varepsilon(t) := & \frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} f_\varepsilon |v - u_1 - \varepsilon u_2|^2 dv dx + \frac{1}{2} \varepsilon \int_{\mathbb{R}} |\partial_x \phi_\varepsilon - \partial_x \phi_1 - \varepsilon \partial_x \phi_2|^2 dx \\ & + \frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} \left( \phi_\varepsilon - \frac{1}{\varepsilon} - \phi_1 - \varepsilon \phi_2 \right)^2 dx. \end{aligned} \quad (5.3)$$

**Acknowledgements:** The author would like to thank David Lannes for stimulating discussions on the long wave limit during the preparation of this work.

## References

1. T. Allemand. The incompressible Euler limit of the Boltzmann equation for a gas of fermions. *Arxiv preprint arXiv:1006.4048*, 2010.
2. B. Alvarez-Samaniego and D. Lannes. Large time existence for 3D water-waves and asymptotics. *Invent. Math.*, 171(3):485–541, 2008.
3. A. A. Arsen'ev. Existence in the large of a weak solution of Vlasov's system of equations. *Ž. Vyčisl. Mat. i Mat. Fiz.*, 15:136–147, 276, 1975.
4. W. Ben Youssef and T. Colin. Rigorous derivation of Korteweg-de Vries-type systems from a general class of nonlinear hyperbolic systems. *M2AN Math. Model. Numer. Anal.*, 34(4):873–911, 2000.
5. W. Ben Youssef and D. Lannes. The long wave limit for a general class of 2D quasilinear hyperbolic problems. *Comm. Partial Differential Equations*, 27(5-6):979–1020, 2002.
6. F. Béthuel, P. Gravejat, J.-C. Saut, and D. Smets. On the Korteweg-de Vries long-wave approximation of the Gross-Pitaevskii equation. I. *Int. Math. Res. Not. IMRN*, (14):2700–2748, 2009.
7. F. Béthuel, P. Gravejat, J.-C. Saut, and D. Smets. On the Korteweg-de Vries long-wave approximation of the Gross-Pitaevskii equation II. *Comm. Partial Differential Equations*, 35(1):113–164, 2010.
8. J. L. Bona, T. Colin, and D. Lannes. Long wave approximations for water waves. *Arch. Ration. Mech. Anal.*, 178(3):373–410, 2005.
9. F. Bouchut. Global weak solution of the Vlasov-Poisson system for small electrons mass. *Comm. Partial Differential Equations*, 16:1337–1365, 1991.
10. F. Bouchut, F. Golse, and M. Pulvirenti. *Kinetic Equations and Asymptotic Theory*. Series in Applied Mathematics. Gauthier-Villars, 2000.
11. J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. II. The KdV-equation. *Geom. Funct. Anal.*, 3(3):209–262, 1993.
12. J. Bourgain. On the Cauchy problem for the Kadomtsev-Petviashvili equation. *Geom. Funct. Anal.*, 3(4):315–341, 1993.
13. Y. Brenier. Convergence of the Vlasov-Poisson system to the incompressible Euler equations. *Comm. Partial Differential Equations*, 25:737–754, 2000.
14. D. Chiron and F. Rousset. The KdV/KP-I limit of the nonlinear Schrödinger equation. *SIAM J. Math. Anal.*, 42(1):64–96, 2010.

15. S. Cordier and Y.J. Peng. Système Euler-Poisson non linéaire. Existence globale de solutions faibles entropiques. *M2AN*, 32(1):1–23, 1998.
16. W. Craig. An existence theory for water waves and the Boussinesq and Korteweg-de Vries scaling limits. *Comm. Partial Differential Equations*, 10(8):787–1003, 1985.
17. V. Duchêne. Boussinesq/Boussinesq systems for internal waves with a free surface, and the KdV approximation. *ESAIM Math. Model. Numer. Anal.*, 46(1):145–185, 2012.
18. V. Duchêne. Decoupled and unidirectional asymptotic models for the propagation of internal waves. *Arxiv preprint arXiv:1208.6394*, 2012.
19. A. V. Faminskii. Well-posed initial-boundary value problems for the Zakharov-Kuznetsov equation. *Electron. J. Differential Equations*, (127):1–23, 2008.
20. F. Golse and L. Saint-Raymond. The Vlasov-Poisson system with strong magnetic field in quasineutral regime. *Math. Models Methods Appl. Sci.*, 13(5):661–714, 2003.
21. E. Grenier. Limite quasineutre en dimension 1. In *Journées “Équations aux Dérivées Partielles” (Saint-Jean-de-Monts, 1999)*, pages Exp. No. II, 8. Univ. Nantes, Nantes, 1999.
22. Y. Guo and X. Pu. KdV limit of the Euler-Poisson system. *Arxiv preprint arXiv:1202.1830*, 2012.
23. Y. Guo and W. A. Strauss. Instability of periodic BGK equilibria. *Comm. Pure Appl. Math.*, 48(8):861–894, 1995.
24. Y. Guo and W. A. Strauss. Nonlinear instability of double-humped equilibria. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 12(3):339–352, 1995.
25. Y. Guo and W. A. Strauss. Unstable BGK solitary waves and collisionless shocks. *Comm. Math. Phys.*, 195(2):267–293, 1998.
26. D. Han-Kwan. Quasineutral limit of the Vlasov-Poisson system with massless electrons. *Comm. Partial Differential Equations*, 36(8):1385–1425, 2011.
27. M. Haragus, D. P. Nicholls, and D. H. Sattinger. Solitary wave interactions of the Euler-Poisson equations. *J. Math. Fluid Mech.*, 5(1):92–118, 2003.
28. E.W. Laedke and K.H. Spatschek. Growth rates of bending KdV solitons. *Journal of Plasma Physics*, 28(03), 1982.
29. D. Lannes, F. Linares, and J.C. Saut. The Cauchy problem for the Euler-Poisson system and derivation of the Zakharov-Kuznetsov equation. *Arxiv preprint arXiv:1205.5080*, 2012.
30. Z. Lin. Instability of periodic BGK waves. *Math. Res. Lett.*, 8(4):521–534, 2001.
31. Z. Lin. Nonlinear instability of periodic BGK waves for Vlasov-Poisson system. *Comm. Pure Appl. Math.*, 58(4):505–528, 2005.
32. Z. Lin and C. Zeng. Small BGK waves and nonlinear Landau damping. *Comm. Math. Phys.*, 306(2):291–331, 2011.
33. Z. Lin and C. Zeng. Small BGK waves and nonlinear Landau damping (higher dimensions). *Arxiv preprint arXiv:1106.4368*, 2011.
34. F. Linares and A. Pastor. Local and global well-posedness for the 2D generalized Zakharov-Kuznetsov equation. *J. Funct. Anal.*, 260(4):1060–1085, 2011.
35. F. Linares and J.-C. Saut. The Cauchy problem for the 3D Zakharov-Kuznetsov equation. *Discrete Contin. Dyn. Syst.*, 24(2):547–565, 2009.
36. N. Masmoudi. From Vlasov-Poisson system to the incompressible Euler system. *Comm. Partial Differential Equations*, 26(9-10):1913–1928, 2001.
37. C. Mouhot and C. Villani. On Landau damping. *Acta Math.*, 207(1):29–201, 2011.
38. X. Pu. Dispersive of the Euler-Poisson system in higher dimensions. *Arxiv preprint arXiv:1204.5435*, 2012.
39. F. Ribaud and S. Vento. Well-Posedness Results for the Three-Dimensional Zakharov-Kuznetsov Equation. *SIAM J. Math. Anal.*, 44(4):2289–2304, 2012.
40. L. Saint-Raymond. Convergence of solutions to the Boltzmann equation in the incompressible Euler limit. *Arch. Ration. Mech. Anal.*, 166(1):47–80, 2003.
41. L. Saint-Raymond. *Hydrodynamic limits of the Boltzmann equation*, volume 1971 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009.
42. L. Saint-Raymond. Hydrodynamic limits: some improvements of the relative entropy method. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 26(3):705–744, 2009.
43. G. Schneider and C. E. Wayne. The long-wave limit for the water wave problem. I. The case of zero surface tension. *Comm. Pure Appl. Math.*, 53(12):1475–1535, 2000.
44. H.T. Yau. Relative entropy and hydrodynamics of Ginzburg-Landau. *Lett. Math. Phys.*, 22(1):63–80, 1991.
45. V.E. Zakharov and E.A. Kuznetsov. On three dimensional solitons. *Sov. Phys. JETP.*, 39:285–286, 1974.
46. Y. Zheng and A. Majda. Existence of global weak solutions to one-component Vlasov-Poisson and Fokker-Planck-Poisson systems in one space dimension with measures as initial data. *Comm. Pure Appl. Math.*, 47(10):1365–1401, 1994.